

**Oxidative Stress and Mitochondrial Injury in Chronic Multisymptom Conditions:
From Gulf War Illness to Autism Spectrum Disorder**

Beatrice Alexandra Golomb, MD, PhD

University of California, San Diego

Department of Medicine

Department of Family and Preventive Medicine

Corresponding Author:

Beatrice Golomb, MD, PhD
University of California, San Diego
9500 Gilman Dr #0995
La Jolla, CA 92093
Phone (858) 558-4950 x201
Fax (858) 558-4960
Email bgolomb@ucsd.edu

Running Title: Gulf War and Mitochondria

Keywords: Gulf War; multisymptom illness; mitochondria; cell energetics; oxidative stress, chronic fatigue syndrome, fibromyalgia, irritable bowel syndrome

Acknowledgments: I gratefully thank Hanh Nguyen, Laura Erickson and Sabrina Koperski for producing the figure; and Sabrina Koperski for invaluable research, administrative and editorial assistance. I thank Sabrina Koperski and Levan Atanelov for comments on the manuscript.

Conflict of Interest: The author has no conflicts of interest to declare.

Abbreviations:

AChEi – Acetylcholinesterase inhibitors
ALS – Amyotrophic Lateral Sclerosis
ASD – Autism Spectrum Disorder
CDC – Center for Disease Control
CFS – Chronic Fatigue Syndrome
CMs – Carbamates
DBP – Diastolic Blood Pressure
DEET – N,N-Diethyl-meta-toluamide

FM – Fibromyalgia
GWI – Gulf War Illness
GWV – Gulf War Veterans
IBS – Irritable Bowel Syndrome
MCS – Multiple Chemical Sensitivity
mtDNA – mitochondrial DNA
NO/NOS – Nitric Oxide / Nitric Oxide Synthase
OS – Oxidative Stress

Golomb 2012 – Gulf War Mitochondria

OSMD – Oxidative Stress and Mitochondrial
Dysfunction
Q10 – Coenzyme Q10

RNS – Reactive Nitrogen Species
ROS – Reactive Oxygen Species
SBP – Systolic Blood Pressure

Abstract

Background: Overlapping chronic multisymptom illnesses (**CMI**) include Chronic Fatigue Syndrome (**CFS**), fibromyalgia, irritable bowel syndrome, multiple chemical sensitivity, and Gulf War illness (**GWI**), and subsets of autism spectrum disorder (**ASD**). GWI entails a more circumscribed set of experiences that may provide insights of relevance to overlapping conditions.

Objectives: To consolidate evidence regarding a role for oxidative stress and mitochondrial dysfunction (**OSMD**), as primary mediators in CMI, using GWI as a departure point.

Methods: Exposure relations, character, timecourse and multiplicity of symptoms, and objective correlates of GWI are compared to expectation for OSMD. Objective correlates of OSMD in GWI and overlapping conditions are examined.

Discussion: OSMD is an expected consequence of known GWI exposures; is compatible with symptom characteristics observed; and accords with objective markers and health conditions linked to GWI, extending to autoimmune disease and infection. Emergent triangulating evidence directly supports OSMD in multisymptom “overlap” CMI conditions, with similarities to, and diagnosed at elevated rates in, GWI, suggesting a common role in each.

Conclusions: GWI is compatible with a paradigm by which uncompensated exposure to oxidative/nitrative stressors accompanies and triggers mitochondrial dysfunction, cell energy compromise, and multiple downstream effects such as vulnerability to autoantibodies. This promotes a profile of protean symptoms with variable latency emphasizing but not confined to energy-demanding post-mitotic tissues, according with (and accounting for) known properties of multisystem overlap conditions. This advances understanding of GWI; health conditions attending GWI at elevated rates; and overlap conditions like CFS and ASD, providing prospects

for vulnerability assessment, mitigation of progression, treatment, and future prevention – with implications germane to additive and excessive environmental oxidative stressor exposures in the civilian setting.

Introduction

Chronic multisymptom illnesses (**CMI**), including chronic fatigue syndrome, fibromyalgia, irritable bowel syndrome, multiple chemical sensitivity and Gulf War illness (**GWI**), show strong overlap¹⁻⁴. Autism spectrum disorder (**ASD**), despite little cooccurrence with GWI (due to military self-selection and selection), encompasses multisymptom subsets bearing muscle, gastrointestinal, sleep and other symptoms germane to CMI. These are suggested to reflect a condition arising from similar mechanisms in a different developmental milieu. GWI entails a circumscribed set of experiences that may provide insights of relevance to these overlapping conditions.

One fourth to 1/3 of US and UK personnel deployed to the Persian Gulf in the 1990-1 conflict exhibit GWI, established by excluding the fraction of nondeployed controls meeting the symptom portion of eligibility⁵⁻⁹. The foundation for GWI has been felt to elude physiological explanation^{10, 11}. GWI differs in exposure relations and symptom profiles from other post-war syndromes (e.g. post-traumatic stress disorder). This conflict had many exposures that were new (anthrax vaccine, depleted uranium, permethrin impregnated uniforms), unique or comparatively unique (pyridostigmine bromide nerve agent pretreatment pills, oil fires, flea collar use by some, botulinum toxoid vaccine, nerve agent exposure resulting from munitions depot demolitions), and excessive (organophosphate and carbamate pesticide use).

Oxidative stress (**OS**) and mitochondrial dysfunction (**MD**) (together **OSMD**) are closely intertwined, and each promotes the other. We evaluate characteristics of OSMD and GWI, to show that OSMD may underlie some, much, or most the excess of multi-symptom illness (and neurodegenerative disease) reported in Gulf War veterans (**GWV**)^{12, 13} – and civilians with CMI.

Method

Exposure relations, symptom characteristics, objective marker correlates, and medical conditions in GWI are examined in relation to OSMD. Evidence from OSMD in overlap conditions are brought to bear in understanding GWI, and vice versa.

Results/Review

Gulf War Illness

26-32% of GWV demonstrate chronic multi-symptom health problems and thus have GWI apparently associated with participation^{5, 7}. (That is the fraction bearing such problems, after subtracting what was “expected” based on nondeployed controls.) The excess burden lies not in more persons with single or dual symptoms, or multiple mild symptoms; but in the added fraction bearing 3 or more symptoms of moderate or greater severity⁵. The 1990-1 ground war lasted but four days; relatively few of the ~700,000 deployed US veterans experienced combat or direct combat stressors. Though some affected veterans have psychological symptoms, most do not meet criteria for psychiatric illness, and stress-related exposures lose significance in multivariable models that adjust for chemical/environmental exposures (but not the converse)^{8, 9}.

Dominant Symptoms; Symptom Multiplicity and Heterogeneity

GWV prominently features fatigue, muscle pain and weakness, and central nervous system symptoms (cognition, mood and personality). The CDC case definition requires chronic symptoms in two of three domains of fatigue, cognitive-mood, and musculoskeletal⁷. Kansas criteria require symptoms in three of six domains – fatigue/sleep, pain (muscle-joint), and cognitive-mood-neurological, gastrointestinal, respiratory and skin⁵; these also must be multiple

within the category and/or at least moderately severe, with onset since 1990⁵. Kansas criteria are more specific (fewer non-GWV meet symptom criteria).

Archetypal clinical symptoms in MD also comprise fatigue, muscle and brain symptoms “Encephalomyopathy”¹⁴⁻²⁰ is the classic manifestation of mitochondrial respiratory chain dysfunction. Over 100 distinct mtDNA mutations have been linked to brain and muscle symptoms²¹. Brain and muscle are postmitotic, with high energy demands, providing particular vulnerability^{21-24 70}.

Symptom multiplicity and heterogeneity further typify GWI. Many additional symptoms, though less frequent, also occur at markedly elevated rates^{5, 10, 25, 26}. Other CMI also commonly bear multiple symptoms, with high overlap^{2, 4}.

Such symptom multiplicity^{21, 27} and heterogeneity^{28-37 38} also characterize MD, so much that “A mitochondrial disease should be considered in the event of dysfunction of more than 2 organ systems or processes with high energy requirements”²⁷. The remaining Kansas GWI criteria -- gastrointestinal dysfunction^{16, 36, 37}, pain^{29, 31, 39}, skin^{40, 41}, sleep⁴²⁻⁴⁵, and breathing symptoms are common in MD^{43, 46}, with many further symptoms elevated in both settings. Symptom heterogeneity in MD rests on a) random somatic segregation of mitochondrial (**mt**) DNA mutations^{47, 48} and mitochondrial heteroplasmy⁴⁹⁻⁵¹: different organs bear different *prior* mtDNA mutation loads leading to different vulnerability; b) variable progression of mitochondrial defects: different mutations may *arise* in different tissues, with some more severe and/or conducive to OS that fosters more mtDNA mutations^{19, 52}; and c) clinical threshold effects^{21, 53}: symptoms arise when mtDNA mutations or resulting cell death surpass a threshold. The fraction of mutated mtDNA in an organ, the mutation severity, the energetic demands of the organ and the cell loss determine clinical consequences^{21, 54}. Symptom variability is an *inherent feature of*

MD (Even persons with familial MD may vary vastly in which symptoms are expressed - and in their timecourse of expression⁵⁵). – and an empiric feature of GWI and other CMI¹⁰. (See **Table 1a&b, Supplement Table 2.**)

Symptom latency

It is often presumed that symptoms, to relate to an exposure, must arise during an exposure, and that the exposure must show acute toxicity. (Cancer and neurodegenerative disease, familiar disease conditions rather than symptoms, represent recognized exceptions.) GWI does not behave in this fashion.

[GWI is characterized by variable latency to symptoms following Gulf exposures](#) – with many new symptoms arising well after participation, and 40% of new symptoms reportedly arising more than a year after Gulf service¹¹. [Symptoms in MD are also characterized by variable latency to onset](#). MD produces (commonly) OS. OS trigger further mtDNA mutations and more MD. The severity of new mutations, and degree of heteroplasmy, determine the severity of OS production – and speed of progression. As mtDNA mutations and OS accrue, cells lose function or die, from energy depletion or OS-precipitated necrosis or apoptosis⁵⁶ – achieving phenotypic thresholds^{21, 53}. (See below for autoimmune predisposition.) Even in a kindred with heritable mtDNA defects that involved neurological problems, “the age of onset of major neurological disturbance varied from 3-70 years”⁵⁵. The principles are similar whether initial mtDNA damage is heritable or acquired; however acquired mtDNA mutations may be typified by multiple different mutations each in a low fraction^{19, 49}, which may particularly dispose to classical or “nonspecific” symptoms of MD, compatible with CMI. OS may predispose to autoantibodies and vulnerability to infection (below). Delayed symptoms may further arise from OS-promoted

autoantibodies and colonization/infection with pathogens (below); and as new OS exposures appear, to which existing OSMD provide heightened vulnerability (due to adverse OS:antioxidant defense balance). OS may promote impaired gene expression related to detoxification, in part via alterations in DNA methylation.**Add excitotoxicity?**

Exposure Associations

A range of “unrelated” exposures are linked to GWI. Acetylcholinesterase inhibitor (AChEi) exposures (organophosphates, as nerve agents and pesticides; and carbamates, as pesticides and pyridostigmine bromide nerve agent pretreatment pills) show especially strong and consistent links to health problems in GWV⁵⁷. Moreover, AChEis show a dose-response relationship to GWI (number of pyridostigmine bromide pills⁵⁷); and proximity to the Khamisiyah munitions depot demolition (sarin nerve agent plume) is linked to extent of brain atrophy and neuropsychological dysfunction in GWV^{58, 59}. Additional findings, extending to genetic evidence (such as paraoxonase variants⁶⁰. These support a causal association of AChEi to GWI⁵⁷, so mechanisms of toxicity by AChEis are of special interest. (GWV/GWI is also linked to reduced paraoxonase activity levels^{60, 61}; however low activity may be effect and/or cause, as paraoxonase is HDL-associated, and OS promotes reductions in HDL.) CMI including chronic fatiguing illness and autism have also been linked to organophosphates⁶²⁻⁶⁴, gene variants in paraoxonase^{62, 65}; and other elements of organophosphate detoxification⁶⁶, as well as low paraoxonase activity⁶⁷.

While AChEi show especially strong and consistent associations to GWI (and are also linked to other CMI), anthrax vaccine^{6, 68-70}, multiple vaccinations^{6, 68, 71-73}, and adverse reactions to vaccinations^{6, 68, 69} show generally consistent significant associations to GWI. Vaccinations are

also linked to other CMI^{64, 74, 75}. Among Gulf-era (Aug 1990-Jul 1991) non-Gulf deployed personnel, rates of “GWI” for those receiving military vaccines in that period OR 3.8 (95%CI 1.5-9.5) were intermediate between rates in non-deployed non-vaccinated (OR 1.0, the standard), and Gulf deployed (OR 10.6, 95% CI 4.9-23.1)⁷⁶ suggesting that vaccines bore a relation to illness irrespective of the Gulf setting. Depleted uranium (bearing potential heavy metal and radioactive toxicity), paints, solvents, and fumes have also shown connections to illness in some epidemiological studies^{5, 6, 26, 69, 71}. Additionally, pesticide exposures of non-AChEi classes also occurred.

The exposures linked to GWI appear “unrelated” in chemical structure and classical mechanism of action. However they share in common induction of toxicity via OSMD. Thus, AChEi toxicity and lethality are normally viewed in cholinergic terms. However in fact they are mediated by OSMD (contributing to apoptosis)^{77, 78}: the salience of OSMD in their toxicity is underscored by the relation of their lethality to impaired glutathione mechanisms; and protection from OP lethality via preexposure or immediate postexposure to relevant antioxidants in experimental animal studies⁷⁹⁻⁸¹. Vaccines, in contrast, are not known to inhibit AChE. However these, and other exposures linked to GWI, each exerts toxicity via OSMD (**Table 1a**). Further, the number of such exposures (crossing different classes) also predicted GWI¹¹ – and predicts greater OSMD⁸²⁻⁸⁵. These considerations begin to provide a framework, coupled with individual variabilities, for genesis of some cases of CMI outside the Gulf War setting.

Proposed Mechanism

OS via reactive oxygen species (**ROS**) and also reactive nitrogen species, (**RNS**), (for simplicity of exposition we designate the joint processes as “OS”) damage proteins, lipids, RNA

and DNA – particularly in mitochondria^{86, 87}, impairing mitochondrial energy production. Mitochondria are a leading target of ROS^{88, 89} due to proximity to ROS production (much of which occurs in mitochondria^{88, 90, 91}), such that mtDNA mutate at 10-1000x the rate of nuclear DNA^{92, 93}. Since all mtDNA genes are germane to oxidative phosphorylation⁹⁴⁻⁹⁶, mtDNA damage commonly hampers mitochondrial respiratory chain function, which in turn further impairs cell energy production and often further increases ROS release^{86, 87}. Reduced energy and increased ROS each cause cell (and subcellular) dysfunction and each can induce cell death, by necrosis or apoptosis^{56, 97-100}. Additionally, the further increase in ROS that is a consequence of mtDNA damage can induce further mtDNA injury – advancing a cycle of OSMD, cell energy depletion, cell dysfunction, and potentially cell loss^{86, 87, 101}. (OS also promotes MD by inhibiting mitochondrial import of essential precursor proteins¹⁰².) OS, adversely affecting the balance of OS to antioxidant defense, can increase vulnerability to, and clinical consequences of, new oxidative exposures. When enough mitochondria are dysfunctional, or enough cells dysfunctional or dead, symptoms or organ dysfunction emerge – mitochondrial “threshold effects”. (OS may have further implications via effects on DNA methylation¹⁰³ and excitotoxicity^{104, 105}, which may be magnified in settings of low mt EN production^{106, 107}.) It is reiterated that major Gulf exposures are known to produce toxicity via OSMD producing expectation of cell dysfunction and cell death. (Note that organophosphates further produce mitochondrial and energetic compromise through toxicity to microtubules¹⁰⁸, interfering with mitochondrial biogenesis¹⁰⁹ and transport¹¹⁰. Mitochondria are dynamic rather than static organelles^{111, 112}.)

Downstream Effects - OS-induced Mechanisms

OS also depress vitamin D (vitD), vitD receptor expression and mitochondrial vitD hydroxylase activities^{113, 114}. Low vitD and altered receptor function are linked to risk of autoimmune disease¹¹⁵⁻¹¹⁹. Autoimmune markers are elevated in GWI^{120, 121}, as well as in other CMI^{74, 75, 122}). Vaccines have been linked to illness in GWV^{6, 68-71} and in some cases chronic fatigue. (They are also a politically and scientifically contentious proposed contributor to ASD.) Reactogenic vaccines are a source of OS^{82♦}. Additionally, vaccine adjuvants (based on aluminum – which is an oxidative stressor, as are components of vaccine preservatives and adjuvants) are expressly incorporated for the purpose of enhancing immune/antibody reactions – in principle to the intended administered antigen, but they may also adjuvant native protein and nonprotein substances. Adjuvants associated with vaccines remain resident in the body and may continue to exert adjuvant effects, promoting “autoimmune syndromes induced by adjuvants” (“ASIA”)^{122, 125-128}. Low vitamin D activity (see above) is also linked to increased infection vulnerability¹²⁹⁻¹³⁴. These factors, coupled with EN deficits and OS-reduced NK activity^{120, 135}, may account for increased autoantibodies, and increased evidence of a range of infectious agents (and antibodies to them), in GWI and associated exposures^{120 136-139} – and other CMI^{74, 75, 140} including ASD¹⁴¹⁻¹⁴³. These contribute further to symptom heterogeneity and latency.

Excitotoxicity is another potential downstream mechanism. Oxidative stress enhances excitotoxin effects, such as DCD (delayed calcium deregulation) which “precedes and predicts” cell death¹⁴⁴. So, too, does mitochondrial calcium accumulation¹⁴⁴. Moreover, excitotoxins in

♦ Squalene has been politically contentious in GWI. One is not impelled to suppose that squalene-adjuvanted variants of anthrax vaccine were used, in order for squalene antibodies to arise: squalene occurs naturally in the body and may also be present in small amounts in vaccines as a contaminant¹²³. McDonald T. MOD Anthrax Vaccine Contains Squalene. "Tonight with Trevor McDonald", March 17, 2003, ITV1, 8PM 2003 Mar 17, 2003, 124. Rodriguez PM. Squalene presence confirmed by FDA. Insight 2000; <http://www.insightmag.com/news/2000/10/30/TheLastWord/Squalene.Presence.Confirmed.By.Fda-213345.shtml>.

turn cause oxidative stress and mitochondrial impairment, which is a major mediator of excitotoxin neurotoxicity¹⁴⁵.

Other sequelae of oxidative stress and mitochondrial dysfunction further contribute to cell dysfunction, cell death, and symptoms.

Differential vulnerability

Differential vulnerability – in which some but not other servicepersons with apparently similar exposures have developed GWI (or CMI) – emerges naturally from an OSMD conceptualization. Differences in genetic and nongenetic biological detoxification capability^{60, 66, 106, 146-150}, total OS load and pro-oxidant/antioxidant balance¹⁴⁷, prior inherited or acquired mitochondrial mutation load or heteroplasmy status^{47, 49, 54, 95}, heritable vulnerability to autoimmune disease, and perhaps prior loss of cells in postmitotic organs, provide for individual differences in vulnerability to development of symptoms following Gulf- (or civilian-) associated OS exposures.

Objective findings

Routine laboratory tests are not classically abnormal in GWV (or CMI) or in OSMD. **Table 1a&b** depict those objective findings that have been reproducibly documented in GWV (demonstrated in at least two studies). In each case the objective finding has a demonstrated relation to OSMD (generally OS, which is more commonly tested). These include depressed paraoxonase (PON1)^{60, 61, 151}, increased gamma glutamyl transferase (GGT)^{152, 153}, reduced natural killer cell activity^{120, 154}, blunted heart rate variability,^{155, 156} increased autoantibodies^{120,}

¹²¹, and increased coagulation activation^{157, 158}. Each has been reported in other CMI. And each is a documented consequence of OSMD.

Associated Health Conditions

In addition to symptoms, several conditions have been found to be elevated in GWV. These include hearing loss, hypertension, and amyotrophic lateral sclerosis (**Table 1a&b**). These conditions are also known to bear particularly strong relations to OSMD.

These findings are compatible with a common mechanism involving OSMD, targeting different tissues to varying degrees. Motor neurons are one possible (though relatively rare) target tissue, engendering ALS when affected. However other distributions of effect result in multisymptom conditions (GWI-CFS-spectrum conditions), metabolic syndrome features¹⁵⁹, and a range of other conditions, which accompany one another at elevated rates^{2-4, 160}.

Triangulating Evidence from CMI

Patients with other CMI commonly have multiple “unexplained” symptoms spanning many domains. These conditions have substantial overlap with one another -- and with GWI^{2-4, 6, 26, 160-164} (see **Table 1a&b**) consistent with the hypothesis put forth here that common OSMD mechanisms target multiple domains, with factors like heteroplasmy (among others) producing differential vulnerability of potential target tissues.

Reported odds ratios for CFS range from ~4 to ~40 in GWV vs nondeployed controls (**Table 1a&b**), suggesting shared mechanisms at least for subsets of CFS. CFS patients show elevated OS¹⁶⁵, an exaggerated prooxidant response to new OS¹⁶⁶ (which we propose may signify a preexisting adverse prooxidant:antioxidant balance), and activation of coagulation

pathways¹⁵⁷ (the relation to oxidative stress is discussed above). Moreover, OS levels correlate with symptoms in CFS¹⁶⁷. Evidence also favors a relation of CFS to MD: functional status in CFS correlates with carnitine levels^{168, 169}; and a subset of CFS cases show elevated lactate with exertion, and delayed ATP recovery after exercise, on magnetic resonance spectroscopy¹⁷⁰, with OS believed to underlie the energetic effects¹⁷¹. Furthermore, an “ATP profile test” of mitochondrial function successfully (and completely) discriminated CFS cases, defined by the CDC criteria which require multisymptom illness (most similar to GWI)¹⁷². Additionally, CFS patients have been reported to differ from controls in genetics related to mitochondrial function¹⁷³, energetics¹⁷³ and apoptosis¹⁷⁴. Fibromyalgia and irritable bowel syndrome²⁶ – which are increased (~2-5 fold) in GWV – also bear evidence of a link to OSMD^{39, 175-177} (**Table 1a&b**). So does ASD^{67, 103, 148, 149, 178-182}. MCS has been linked to mutations in genes for glutathione-S-transferase (GST), which protects from oxidative stress by conjugating glutathione, detoxifying “a large range of compounds generated by reactive oxygen species induced damage to intracellular molecules”.¹⁴⁷ It might be conjectured that apparent instances of exposure-triggered MCS may entail exposure-triggered increased endogenous OS production (through mechanisms detailed above); and perhaps also exposure-triggered modifications in gene expression of key detoxifying enzymes.

MCS, “sensitivity” to chemicals – and also medication intolerance¹⁸³ – are significantly more common in GWV^{5, 6, 25, 26, 160, 184, 185} and in overlap conditions^{2-4, 186}. Both chemical (**Table 3**) and drug adverse effects are commonly mediated through OSMD¹⁸⁷⁻¹⁹⁰. (MCS, like GWI and ASD cited elsewhere, shows altered genetic profiles of PON^{60, 191}.) OSMD provides an account of why medication and vaccine adverse effects at the time of the Gulf were associated with increased risk of developing GWI^{6, 68, 192, 193}. These adverse effects signal less favorable OSMD status at the time of exposure *and or* greater oxidative

stressor exposure, promoting OS dominance over antioxidant defences, which may both predict and help mediate the development of multisymptom GWI in the setting of new OS exposure.

Other conditions have been linked to OSMD but have not (yet) been evaluated in GWI. An elevation in a number of these conditions in GWI represents a prediction of this theory. Some markers and objective conditions have been linked to GWI and OSMD, and at least one CMI; but have not been evaluated in other CMI. A prediction of this theory is that many, when examined in studies with sufficient power, will be seen in other CMI.

Discussion

An explanation focused on OSMD accounts for GWI, and other CMI. Put forth is a proposition by which exposures to oxidative stressors, in part via cumulative MD, in part via autoimmune effects in vulnerable individuals, in part via apoptosis/cell loss leading to coagulation activation, may increase risk of a range of chronic health problems in GWV – including the multisymptom spectrum termed GWI – and other CMI. Such an account accords with known exposure relations of otherwise unlike character; explains symptoms with otherwise unexplained organ tropism, with otherwise unexplained protean character and with otherwise unexplained variable timecourse to onset. It fits with the otherwise unexplained range of objective markers linked to these conditions that cross classical boundaries. It explains observed associations to other health conditions of varying character; and accounts for increases in multisymptom “overlap” conditions (and for overlap of these syndromes with one another). A role for OSMD might be predicted on “first principles” (known pathophysiological effects of Gulf Exposures; and known clinical sequelae of these effects)¹³. However there is now direct evidence of a relation of OS and MD to each of the conditions to which GWI relates, providing further triangulating support for a role for OS and MD extending to GWI.

CMI has not traditionally been thought to include autism spectrum disorder (**ASD**); but multisymptom cases of ASD may reflect manifestations of similar processes operating at a different developmental timepoint. ASD not uncommonly comprises multiple symptoms across a similar spectrum,^{181, 194-197} with purported links to similar exposures, and observed relations to OSMD^{106, 149, 180, 182, 198-207}. (All ASD need not cohere with these principles.)

Accrued evidence now more vigorously supports a paradigm by which OS and MD conduce to one another, disposing to GWI or other CMI. These mechanisms may operate in some, many or possibly most CMI. The Gulf War setting has its distinctive constellation of OS exposures. However the pathophysiological principles articulated, beyond their potential relevance to GWI, yield a new lens with which to review a range of important, interrelated health conditions – including but not confined to “overlap” CMI conditions often considered to be unexplained – and to understand their relationship to one another – in the military setting, and outside of it.

Figure 1: Oxidative Stress and Mitochondrial Dysfunction Mediate the Link between “Unrelated” Exposures and Symptoms




Table 1. Overview

	GW	OSMD	CMI
Dominant Symptoms	Symptoms protean but focus on fatigue, cognitive-mood, musculoskeletal, also gastrointestinal, sleep, neurological	Symptoms protean but focus for MD on fatigue, cognitive-mood, musculoskeletal, also gastrointestinal, sleep, neurological	Symptoms protean with emphasis defined by the condition (fatigue for CFS; muscle for FM; GI for IBS; CNS/"cognitive" for ASD; chemical sensitivity for MCS) but very high overlap
Symptom multiplicity/heterogeneity	Symptoms multiplicity with high heterogeneity	Symptoms multiplicity with high heterogeneity	High heterogeneity of symptoms
Symptom Latency	Variable Latency to Symptom Onset	Variable Latency to Symptom Onset	Variable Latency to Symptom Onset
Exposure Associations	Include OPs/ acetylcholinesterase inhibitors, paraoxonase gene variants, vaccines	Causes include OPs/acetylcholinesterase inhibitors, paraoxonase gene variants, vaccines	Include OPs/ acetylcholinesterase inhibitors, paraoxonase gene variants, vaccines
Objective Findings	Include autonomic dysfunction, reduced natural killer cell activity, coagulation	Include autonomic dysfunction, reduced natural killer cell activity, coagulation	Include autonomic dysfunction, reduced natural killer cell activity ^{208, 209} ,

Golomb 2012 – Gulf War Mitochondria

	activation, elevated autoimmune markers, elevated GGT, low paraoxonase activity	activation, elevated autoimmune markers, elevated GGT, low paraoxonase activity	coagulation activation ²¹⁰ , elevated autoimmune markers ⁷⁴ , ^{75, 122} , elevated GGT ²¹¹ , low paraoxonase activity ⁶⁷
Related conditions	Include CMI (except ASD). Also include hypertension, hearing loss, ALS	Include CMI (extending to ASD). Also include hypertension, hearing loss, ALS	Include hearing loss.

Table 1a. Gulf War Associations

Associated Characteristic	Seen With	
	Gulf War Illness (also other CMI)	Oxidative stress \longleftrightarrow mt Dysfunction
Exposure Associations		
Acetylcholinesterase inhibitors strongly associated	Yes ^{6, 57}	Yes ^{77, 78}
Reactogenic vaccines also consistently associated	Anthrax vaccine associated ^{6, 69} Vaccine adverse effects associated ^{6, 193} Multiple vaccinations associated ^{6, 68, 71, 72}	Reactogenic vaccines associated with OS ⁸² ; Anthrax vaccine among the most reactogenic of vaccines ²¹² Aluminum (vaccine adjuvants), mercury (thimerosal preservatives) linked OS, impaired antioxidant defense against other exposures, and MD ²¹³⁻²²⁸
Multiple other exposures are associated	See Supplement Table 1	1. See Supplement Table 1
Symptom Characteristics		
Brain and muscle dominate	Yes ⁷	Yes ²²
Fatigue and fatigue with exertion prominent	Yes ^{5, 7, 25, 26}	Yes ^{46, 172, 229, 230}
Symptoms are protean spanning many domains	Yes ^{5, 25, 26}	Yes ²³¹
Symptoms differ from person to person	Yes ^{5, 7, 25, 26}	Yes ^{55, 232-234}
Latency to symptom onset is variable, often prolonged, differs across individuals and differs across symptoms within an individual	Yes ¹¹	Yes ⁵⁵
Individual Symptoms Associated	See Supplement Table 2	See Supplement Table 2
Objective Findings		
Routine labs generally unremarkable	Yes	Yes
PON genotype differences present	Yes*** Other CMI ¹⁹¹ add ASD, CFS	Yes
GGT elevated (marker of oxidative stress)	Yes ^{152, 153} (other OS markers not examined) Other CMI:	Yes ^{235, 236}
PON1 activity reduced	Yes ^{60, 61, 151} Other CMI:	Yes ^{237, 238}
Natural killer cell activity reduced	Yes ^{120, 154} Other CMI:	Yes ^{135, 239-241}
Heart Rate variability blunted	Yes ^{155, 156}	Yes ^{242, 243}
Other autonomic abnormalities present	Yes ^{244, 245} Other CMI:	Yes ^{31, 246, 247}
Coagulation markers elevated	Yes ^{157, 158} Other CMI:	Yes ²⁴⁸⁻²⁵¹ (OS promotes apoptosis ^{98, 99, 252} ; which activates coagulation pathways ²⁵³)
Autoimmune Markers Elevated	Yes ^{120, 121} Other CMI: ^{74, 75, 210} autism autoantibodies check	Yes ADD CITS ²⁵⁴ (May be partly adaptive ^{255, 256})

Exposures:

AChEi are particularly potent in inducing OS and MD⁷⁷ – and particularly strong consistent predictors of GWI. AChEi exposures, moreover, were also commonly recurrent when they occurred; and in addition, they cause MD by means beyond OS (such as microtubule toxicity^{257, 258}). The central role of OS in mediating AChEi

Golomb 2012 – Gulf War Mitochondria

toxicity is underscored by findings that pretreatment (or immediate posttreatment) with antioxidants protects against AChEi-induced lethality (and enhances recovery) in animals⁷⁹⁻⁸¹. Anthrax vaccinations and multiple vaccinations are the next most strongly and consistently linked to GWI, in epidemiologic evidence²⁵⁹. These are tied, through aluminum-based adjuvants^{218, 219, 222-224, 226, 260, 261} and mercury (and other) preservatives²²⁵, to OS and MD (**Table 1, Supplement Table 1**) – as well as to inhibition of glutathione production, reducing antioxidant defense against other exposures^{219, 222, 225}. Each additional exposure class presumptively inculcated in GWI is known to promote OS^{82-85, 226, 262-264}.

Markers:

GGT is an enzymatic antioxidant^{265, 266}: elevations serve as an early marker of OS²⁶⁷ including AChEi exposure^{265, 268}. PON1 is an antioxidant enzyme (also specifically involved in organophosphate detoxification²⁶⁹) whose activity is depressed in settings of elevated OS²³⁷. PON1 was depressed in GWV generally (relative to military controls) – and particularly so in those with GWI or more symptoms^{60, 61, 151}. Natural killer activity is reduced with OS and increased with antioxidants^{135, 239, 240}. Blunted heart rate variability has been linked to OSMD^{242, 243}. Activation of the coagulation system relates to OS^{248, 249, 251, 270}. Both OS^{98, 99, 252} and MD^{271, 272} each foster apoptosis; which exposes phosphatidylserine at the cell surface, activating coagulation pathways²⁵³. Activation of the coagulation system has been repeatedly observed in Gulf War veterans.^{157, 158} OS also depresses vitamin D (vitD), vitD receptor expression and mitochondrial vitD hydroxylase activities^{113, 114}. Low vitD and altered receptor function are linked to risk of autoimmune disease¹¹⁵⁻¹¹⁹.

OS – oxidative stress; GWV – gulf war veterans; GWI – gulf war illness; MD – mitochondrial dysfunction

Table 1b. Associated Conditions are Compatible with OSMD

Associated Conditions	Linked to GWV	Linked to OSMD
Chronic Fatigue syndrome	Yes ^{5, 6, 26, 160, 164, 273}	Yes ^{165-167, 170-172, 274-276}
Fibromyalgia	Yes ^{5, 164, 273}	Yes ^{39, 175-177, 277, 278}
Irritable Bowel syndrome	Yes ²⁶	Yes ²⁷⁹
Multiple Chemical Sensitivity	Yes ^{5, 6, 25, 26, 160, 184, 185}	Yes ¹⁴⁷
<i>Autism Spectrum Condition</i>	<i>No – different developmental timeline</i>	Yes ^{67, 106, 107, 148, 180-182, 200, 280, 281}
Hypertension	Yes ^{6, 26, 153, 282, 283}	Yes ²⁸⁴⁻²⁸⁹
Weight gain	Yes ^{7, 26, 153}	Yes ²⁹⁰⁻²⁹³
Amyotrophic lateral sclerosis	Yes ²⁹⁴⁻²⁹⁶	Yes ²⁹⁷⁻³⁰⁰
Hearing loss / tinnitus	Yes ^{5, 6, 10, 26}	Yes ^{28, 234, 301-307}
Fracture risk and reduced bone formation	Yes ^{308, 309}	Yes ³¹⁰⁻³¹³

Significant elevations in self-reported hearing loss after 1990 or 1991 in GWV vs control groups have been reported^{5, 6, 10, 26}. (However obviously, even with military controls, there are uncertainties regarding noise exposure comparability. Additionally, formal testing has not yet been undertaken.) Hearing loss deemed “age-related” and “noise-related” is powerfully linked to OSMD³⁰³⁻³⁰⁶. Accelerated hearing loss is common in MD^{28, 34, 314-322}, cited as among the most common symptoms of mitochondriopathy, together with fatigue and muscle symptoms³⁰. And chemical exposures have been linked to both hair cell damage and *noise*-related hearing loss^{323, 324}.

GWV demonstrate increased incident hypertension in a range of studies (**Table 1a&b**)^{6, 26, 153, 282}. Hypertension, too, is a condition with powerful relationship to OSMD^{284-287, 325, 326}. Indeed an evaluation of mitochondrial pedigrees estimated the fraction of patients with hypertension potentially due to mtDNA mutation involvement at 55% (95% CI 45-65%)²⁸⁵. Analogous to the Gulf War situation, increased blood pressure has been previously associated with environmental OS exposures. For instance, both arsenic and lead are reported to mediate their toxicity via OS³²⁷⁻³²⁹; and exposure to each arsenic (e.g. in the water supply)^{330, 331} and lead^{327, 328} have been linked to increased rates of hypertension. Hypertension and hearing loss are both common “age-related” conditions.

In contrast, amyotrophic lateral sclerosis (**ALS**) is an uncommon but serious age-related condition. ALS has been reported in several studies to have increased incidence in GWV (**Table 1a&b**). This condition has been conceptualized as resulting from “oxidative damage to mitochondrial DNA leading to the accumulation of mitochondrial DNA mutations”³³² – in this case targeting the spinal cord – compatible with the mechanism I propose for GWI more generally – and other GWV associated conditions. Extensive evidence links ALS to OSMD^{297, 299, 300, 333-343}. Moreover, low concentrations and mutations of paraoxonase, central to organophosphate (AChEi) detoxification, have been linked to ALS³⁴⁴⁻³⁴⁷ – and to GWI^{60, 61, 151, 348} (as well as to GWI-like multisymptom illness following organophosphate exposure outside the Gulf setting^{62, 349, 350}), further

* (Noise “sensitivity” has been noted as a some-time feature of multiple chemical sensitivity, and is also noted in ASD. Since both conditions have been linked to impaired antioxidant defenses¹⁴⁷. Schnakenberg E, Fabig KR, Stanulla M, et al. A cross-sectional study of self-reported chemical-related sensitivity is associated with gene variants of drug-metabolizing enzymes. *Environ Health* 2007;6:6, 180. Geier DA, Kern JK, Garver CR, et al. Biomarkers of environmental toxicity and susceptibility in autism. *J Neurol Sci* 2008, 198. Jill James S, Melnyk S, Jernigan S, Hubanks A, Rose S, Gaylor DW. Abnormal Transmethylation/transsulfuration Metabolism and DNA Hypomethylation Among Parents of Children with Autism. *J Autism Dev Disord* 2008, 280. Bradstreet JJ, Smith S, Baral M, Rossignol DA. Biomarker-guided interventions of clinically relevant conditions associated with autism spectrum disorders and attention deficit hyperactivity disorder. *Altern Med Rev* 2010;15:15-32. this “sensitivity” to both noise and chemicals could signify actual vulnerability to damage at lower levels than would be harmful in others.)

Golomb 2012 – Gulf War Mitochondria

propelling the case for OSMD in GWI. By the mechanisms elucidated here (and previously elaborated by us in a different setting of exposure-induced ALS³⁵¹), although spinal cord motor neurons are the primary condition-defining target, mitochondrial dysfunction in sporadic ALS *should not be confined to spinal cord motor neurons* but should (statistically) be evident in other tissues, particularly energetically demanding (and thus mitochondrially vulnerable) brain and muscle. Indeed, recent studies show that mitochondrial pathology in ALS extends to skeletal muscle³⁵²⁻³⁵⁷ and symptoms extend to cognitive function in about half of cases^{358, 359}.

Mounting evidence supports a connection of OSMD^{290-293, 360, 361} to metabolic syndrome (each has been conceptualized as a unifying mechanism^{293, 361}), of which hypertension is one component. Consistent with this, studies have also reported increased incident weight gain in GWV vs controls^{7, 26, 153}. Assessment for other elements of metabolic syndrome in GWV has not yet been undertaken, but increased metabolic syndrome represents a prediction of this model.

Table 3. Predictions

	Observed with Oxidative Stress/ Mitochondrial Dysfunction	Preliminary Evidence or Comment
Other markers of oxidative stress and mitochondrial dysfunction will be elevated	Yes ⁵⁴	True for chronic fatigue syndrome ^{167, 171, 172} , a related condition that is elevated in GWV ^{5, 6, 26} True for ASD
Genetic risk markers related to mitochondrial function, antioxidant defense, energetics, and apoptosis will be identified	Yes, by definition	Gene associations related to mitochondrial function, energetics and apoptosis have been reported for chronic fatigue syndrome ^{66, 173, 174, 276} , a related condition that is elevated in GWV ^{5, 6, 26} . Deletions in genes for glutathione-S-transferase enzymes (leading to loss of protection against oxidative stress) are strongly and significantly linked to multiple chemical sensitivity ¹⁴⁷ Gene variants in paraoxonase have been observed in both GWI and ASD**, and glutathione-related variants among others in ASD ^{148, 198, 362}
GWV will have increased rates of adverse effects with many medications and interventions because many adverse effects are mediated via oxidative stress and mitochondrial problems ¹⁸⁷⁻¹⁹⁰	See Comment. Examples of drugs with adverse effects mediated by oxidative stress include acetaminophen, aminoglycosides, fluoroquinolones, amiodarone, nonsteroidal anti-inflammatory agents, chemotherapeutics, ^{83, 188, 190, 363-378}	Increased drug adverse effects have been observed in irritable bowel syndrome ³⁷⁹ – a condition elevated in GWV ²⁶ and linked to oxidative stress ²⁷⁹ . Regression reported by some parents following vaccinations in cases of ASD may be a <i>consequence</i> of impaired detoxification. (Such vaccination may boost the oxidative burden, however, and potentiate the process.)
New oxidative stressor exposures – beyond medications and medical interventions, though those merit separate mention - will have disproportionate adverse impact in exposed and particularly symptomatic GWV and CMI		Prooxidant-antioxidant balance will be adversely affected, disadvantaging ability to defend against new oxidative stressors.
Endothelial dysfunction will arise at elevated rates	Endothelial dysfunction is strongly linked to oxidative stress ³⁸⁰	
Hepatic steatosis will develop at elevated rates, as will ectopic fat deposition in other locations	Hepatic steatosis relates strongly to oxidative stress ^{381, 382} .	
<i>Low HDL and high triglycerides will emerge</i>	<i>These conditions (and all elements of metabolic syndrome) are linked to oxidative stress^{290, 383} and mitochondrial dysfunction²⁹²</i>	
<i>Free fatty acids will be elevated</i>	<i>Elevated free fatty acids are linked to oxidative stress³⁸⁴</i>	
Metabolic syndrome as a whole will arise at elevated rates	Metabolic syndrome is linked to oxidative stress ^{290, 291} and mitochondrial dysfunction ²⁹²	As above, weight gain and hypertension are already reported at elevated rates. Diabetes prevalence shows variable trends likely due to selection of diabetics out of deployment to high risk areas.
Rates of diabetes will rise to first match then exceed rates in nondeployed (those with diabetes were disproportionately nondeployed)	Diabetes, like metabolic syndrome, is strongly linked to oxidative stress (as effect but possibly also cause) ³⁸⁵ and mitochondrial dysfunction ^{287, 386}	This prediction is made despite mixed direction of trends in diabetes rates in published studies ^{5, 26} . It is predicted that a clear increase will emerge.
Rates of cardiovascular disease and particularly peripheral arterial disease will emerge at an elevated rate as GWV age	These conditions, and especially peripheral arterial disease, are associated with oxidative stress ³⁸⁷⁻³⁸⁹ and mitochondrial dysfunction ³⁹⁰⁻³⁹²	
Parkinson's disease will emerge at an elevated rate as GWV age	Parkinson's disease is linked to oxidative stress and mitochondrial dysfunction ^{393, 394}	Like ALS (and Gulf War illness), Parkinson's disease bears an association to polymorphisms in paraoxonase; and exposure to pesticides ³⁹⁵⁻³⁹⁹
Elevated "depression" diagnoses will be found to disproportionately focus on somatic symptoms	Somatic symptoms in depression have been found to serve as markers for mitochondrial dysfunction. ⁴⁰⁰	"Depression" has been reported to be elevated in GWV ²⁶ . GWV have many somatic symptoms that will contribute to elevated scores in depression scales.
"Age-related" hearing loss and tinnitus will be demonstrated to occur at elevated rates	302, 306, 314, 323, 324	GWV report physician-diagnosed tinnitus at elevated rates ²⁶
Other diagnoses linked to oxidative stress will occur at elevated rates		Some reports suggest increased cancer of some types ²⁸² and cancer death in Gulf War

Golomb 2012 – Gulf War Mitochondria

		veterans ^{401, 402}
<i>Exposed</i> GWV who do not meet criteria for GWI will nonetheless show statistical abnormalities in markers of oxidative stress – and sequelae of oxidative stress (perhaps intermediate between unexposed and those with GWI)		Asymptomatic persons with mt dysfunction at low heteroplasmy rates nonetheless show increased oxidative stress, proportional to the heteroplasmy level ⁵⁴ .

GWI – gulf war illness; GWV – gulf war veterans; ALS – Amyotrophic lateral sclerosis;

* Those with GWI have many somatic symptoms that will contribute to elevated scores in depression scales.

References

1. Frissora CL, Koch KL. Symptom overlap and comorbidity of irritable bowel syndrome with other conditions. *Curr Gastroenterol Rep* 2005;7:264-71.
2. Buchwald D, Garrity D. Comparison of patients with chronic fatigue syndrome, fibromyalgia, and multiple chemical sensitivities. *Arch Intern Med* 1994;154:2049-53.
3. Brown MM, Jason LA. Functioning in individuals with chronic fatigue syndrome: increased impairment with co-occurring multiple chemical sensitivity and fibromyalgia. *Dyn Med* 2007;6:6.
4. Aaron LA, Buchwald D. A review of the evidence for overlap among unexplained clinical conditions. *Ann Intern Med* 2001;134:868-81.
5. Steele L. Prevalence and patterns of Gulf War illness in Kansas veterans: association of symptoms with characteristics of person, place, and time of military service. *Am J Epidemiol* 2000;152:992-1002.
6. Unwin C, Blatchley N, Coker W, et al. Health of UK servicemen who served in Persian Gulf War. *Lancet* 1999;353:169-78.
7. Fukuda K, Nisenbaum R, Stewart G, et al. Chronic multisymptom illness affecting Air Force veterans of the Gulf War. *Jama* 1998;280:981-8.
8. Binns JH, Cherry N, Golomb BA, et al. Research Advisory Committee on Gulf War Veterans' Illnesses: Scientific Progress in Understanding Gulf War Veterans' Illnesses: Report and Recommendations. http://www1.va.gov/rac-gwvi/docs/ReportandRecommendations_2004.pdf; 2004 September 2004.
9. Binns JH, Barlow C, Bloom FE, et al. Gulf War Illness and the Health of Gulf War Veterans. Scientific Findings and Recommendations. Washington D.C.: U.S. Government Printing Office; 2008.
10. Doebbeling BN, Clarke WR, Watson D, et al. Is there a Persian Gulf War syndrome? Evidence from a large population-based survey of veterans and nondeployed controls. *Am J Med* 2000;108:695-704.
11. Kroenke K, Koslowe P, Roy M. Symptoms in 18,495 Persian Gulf War veterans. Latency of onset and lack of association with self-reported exposures. *J Occup Environ Med* 1998;40:520-8.
12. Golomb BA. Mitochondrial dysfunction. A mechanism of illness in Gulf War veterans? Written brief for Dept of Veterans Affairs Research Advisory Committee on Gulf War Illnesses 2002.
13. Golomb BA. Mitochondrial function and Gulf War Illnesses. Presentation to the Department of Veterans Affairs Research Advisory Committee on Gulf War Veterans' Illnesses. Washington DC: Nov 2002. 2002.
14. Boitier E, Degoul F, Desguerre I, et al. A case of mitochondrial encephalomyopathy associated with a muscle coenzyme Q10 deficiency. *Journal of the Neurological Sciences* 1998;156:41-6.
15. Checcarelli N, Prella A, Moggio M, et al. Multiple deletions of mitochondrial DNA in sporadic and atypical cases of encephalomyopathy. *J Neurol Sci* 1994;123:74-9.
16. Coskun E, Ulusal G, Bulut N, Bektas H, Oztekin MF, Yildirim IS. Mitochondrial neurogastrointestinal encephalomyopathy. *Turk J Gastroenterol* 2005;16:163-6.
17. Donovan TJ. Mitochondrial encephalomyopathy: a rare genetic cause of sensorineural hearing loss. *Ann Otol Rhinol Laryngol* 1995;104:786-92.
18. Hara K, Yamamoto M, Anegawa T, Sakuta R, Nakamura M. [Mitochondrial encephalomyopathy associated with parkinsonism and a point mutation in the mitochondrial tRNA(Leu)(UUR) gene]. *Rinsho Shinkeigaku* 1994;34:361-5.
19. Kovalenko SA, Tanaka M, Yoneda M, Iakovlev AF, Ozawa T. Accumulation of somatic nucleotide substitutions in mitochondrial DNA associated with the 3243 A-to-G tRNA(Leu)(UUR) mutation in encephalomyopathy and cardiomyopathy. *Biochem Biophys Res Commun* 1996;222:201-7.
20. Kuhl CK, Layer G, Traber F, Zierz S, Block W, Reiser M. Mitochondrial encephalomyopathy: correlation of P-31 exercise MR spectroscopy with clinical findings. *Radiology* 1994;192:223-30.

21. Wei YH. Mitochondrial DNA mutations and oxidative damage in aging and diseases: an emerging paradigm of gerontology and medicine. *Proc Natl Sci Counc Repub China B* 1998;22:55-67.
22. De Vivo DC, DiMauro S. Mitochondrial defects of brain and muscle. *Biol Neonate* 1990;58 Suppl 1:54-69.
23. Fehm HL, Kern W, Peters A. The selfish brain: competition for energy resources. *Prog Brain Res* 2006;153:129-40.
24. Shulman RG, Rothman DL, Behar KL, Hyder F. Energetic basis of brain activity: implications for neuroimaging. *Trends Neurosci* 2004;27:489-95.
25. Cherry N, Creed F, Silman A, et al. Health and exposures of United Kingdom Gulf war veterans. Part I: The pattern and extent of ill health. *Occup Environ Med* 2001;58:291-8.
26. Gray GC, Reed RJ, Kaiser KS, Smith TC, Gastanaga VM. Self-reported symptoms and medical conditions among 11,868 Gulf War-era veterans: the Seabee Health Study. *Am J Epidemiol* 2002;155:1033-44.
27. Smits BW, Smeitink JA, van Engelen BG. [Mitochondrial diseases; thinking beyond organ specialism necessary]. *Ned Tijdschr Geneesk* 2008;152:2275-81.
28. Chen JC, Tsai TC, Liu CS, Lu CT. Acute hearing loss in a patient with mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like episodes (MELAS). *Acta Neurol Taiwan* 2007;16:168-72.
29. Chu CC, Huang CC, Fang W, Chu NS, Pang CY, Wei YH. Peripheral neuropathy in mitochondrial encephalomyopathies. *Eur Neurol* 1997;37:110-5.
30. Fattal O, Budur K, Vaughan AJ, Franco K. Review of the literature on major mental disorders in adult patients with mitochondrial diseases. *Psychosomatics* 2006;47:1-7.
31. Higashimoto T, Baldwin EE, Gold JI, Boles RG. Reflex sympathetic dystrophy: complex regional pain syndrome type I in children with mitochondrial disease and maternal inheritance. *Arch Dis Child* 2008;93:390-7.
32. Inagaki T, Ishino H, Seno H, Ohguni S, Tanaka J, Kato Y. Psychiatric symptoms in a patient with diabetes mellitus associated with point mutation in mitochondrial DNA. *Biol Psychiatry* 1997;42:1067-9.
33. Shinkai T, Nakashima M, Ohmori O, et al. Coenzyme Q10 improves psychiatric symptoms in adult-onset mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like episodes: a case report. *Aust N Z J Psychiatry* 2000;34:1034-5.
34. Sakaue S, Ohmuro J, Mishina T, et al. A case of diabetes, deafness, cardiomyopathy, and central sleep apnea: novel mitochondrial DNA polymorphisms. *Tohoku J Exp Med* 2002;196:203-11.
35. Tatsumi C, Takahashi M, Yorifuji S, Kitaguchi M, Tarui S. Mitochondrial encephalomyopathy, ataxia, and sleep apnea. *Neurology* 1987;37:1429-30.
36. Aksoy F, Demirel G, Bilgic T, Gungor IG, Ozelik A. A previously diagnosed mitochondrial neurogastrointestinal encephalomyopathy patient presenting with perforated ileal diverticulitis. *Turk J Gastroenterol* 2005;16:228-31.
37. Hirano M, Nishigaki Y, Marti R. Mitochondrial neurogastrointestinal encephalomyopathy (MNGIE): a disease of two genomes. *Neurologist* 2004;10:8-17.
38. DiMauro S, Bonilla E, Lombes A, Shanske S, Minetti C, Moraes CT. Mitochondrial encephalomyopathies. *Neurol Clin* 1990;8:483-506.
39. van de Glind G, de Vries M, Rodenburg R, Hol F, Smeitink J, Morava E. Resting muscle pain as the first clinical symptom in children carrying the MTTK A8344G mutation. *Eur J Paediatr Neurol* 2007;11:243-6.
40. Flynn MK, Wee SA, Lane AT. Skin manifestations of mitochondrial DNA syndromes: case report and review. *J Am Acad Dermatol* 1998;39:819-23.
41. Kira R, Kubo E, Hara T. [Endocrine, skin, and bone marrow involvement in mitochondrial cytopathies; a review]. *Nippon Rinsho* 2002;60 Suppl 4:659-61.
42. Cerveri I, Fanfulla F, Zoia MC, Manni R, Tartara A. Sleep disorders in neuromuscular diseases. *Monaldi Arch Chest Dis* 1993;48:318-21.
43. Manni R, Piccolo G, Banfi P, et al. Respiratory patterns during sleep in mitochondrial myopathies with ophthalmoplegia. *Eur Neurol* 1991;31:12-7.

44. Suzuki Y, Taniyama M, Hata T, Miyaoka H, Atsumi Y, Matsuoka K. Sleep-wake dysrhythm in mitochondrial diabetes mellitus. *Diabetes Res Clin Pract* 1997;35:61-2.
45. Tatsumi C, Takahashi M, Yorifuji S, et al. Mitochondrial encephalomyopathy with sleep apnea. *Eur Neurol* 1988;28:64-9.
46. Flaherty KR, Wald J, Weisman IM, et al. Unexplained exertional limitation: characterization of patients with a mitochondrial myopathy. *Am J Respir Crit Care Med* 2001;164:425-32.
47. Kirches E, Michael M, Warich-Kirches M, et al. Heterogeneous tissue distribution of a mitochondrial DNA polymorphism in heteroplasmic subjects without mitochondrial disorders. *J Med Genet* 2001;38:312-7.
48. Matthews PM, Brown RM, Morten K, Marchington D, Poulton J, Brown G. Intracellular heteroplasmy for disease-associated point mutations in mtDNA: implications for disease expression and evidence for mitotic segregation of heteroplasmic units of mtDNA. *Hum Genet* 1995;96:261-8.
49. Smigrodzki RM, Khan SM. Mitochondrial microheteroplasmy and a theory of aging and age-related disease. *Rejuvenation Res* 2005;8:172-98.
50. Nagley P, Mackay IR, Baumer A, et al. Mitochondrial DNA mutation associated with aging and degenerative disease. *Ann N Y Acad Sci* 1992;673:92-102.
51. Wallace DC, Zheng XX, Lott MT, et al. Familial mitochondrial encephalomyopathy (MERRF): genetic, pathophysiological, and biochemical characterization of a mitochondrial DNA disease. *Cell* 1988;55:601-10.
52. Wei YH. Oxidative stress and mitochondrial DNA mutations in human aging. *Proc Soc Exp Biol Med* 1998;217:53-63.
53. Rossignol R, Faustin B, Rocher C, Malgat M, Mazat JP, Letellier T. Mitochondrial threshold effects. *Biochem J* 2003;370:751-62.
54. Canter JA, Eshaghian A, Fessel J, et al. Degree of heteroplasmy reflects oxidant damage in a large family with the mitochondrial DNA A8344G mutation. *Free Radic Biol Med* 2005;38:678-83.
55. Crimmins D, Morris JG, Walker GL, et al. Mitochondrial encephalomyopathy: variable clinical expression within a single kindred. *J Neurol Neurosurg Psychiatry* 1993;56:900-5.
56. Odinokova IV, Sung KF, Mareninova OA, Hermann K, Gukovsky I, Gukovskaya AS. Mitochondrial mechanisms of death responses in pancreatitis. *J Gastroenterol Hepatol* 2008;23 Suppl 1:S25-30.
57. Golomb BA. Acetylcholinesterase inhibitors and Gulf War illnesses. *Proc Natl Acad Sci U S A* 2008;105:4295-300.
58. Heaton KJ, Palumbo CL, Proctor SP, Killiany RJ, Yurgelun-Todd DA, White RF. Quantitative magnetic resonance brain imaging in US army veterans of the 1991 Gulf War potentially exposed to sarin and cyclosarin. *Neurotoxicology* 2007;28:761-9.
59. Proctor SP, Heaton KJ, Heeren T, White RF. Effects of sarin and cyclosarin exposure during the 1991 Gulf War on neurobehavioral functioning in US army veterans. *Neurotoxicology* 2006;27:931-9.
60. Haley RW, Billecke S, La Du BN. Association of low PON1 type Q (type A) arylesterase activity with neurologic symptom complexes in Gulf War veterans. *Toxicol Appl Pharmacol* 1999;157:227-33.
61. Mackness B, Durrington PN, Mackness MI. Low paraoxonase in Persian Gulf War Veterans self-reporting Gulf War Syndrome. *Biochem Biophys Res Commun* 2000;276:729-33.
62. Cherry N, Mackness M, Durrington P, et al. Paraoxonase (PON1) polymorphisms in farmers attributing ill health to sheep dip. *Lancet* 2002;359:763-4.
63. Davies DR, Ahmed GM, Freer T. Chronic organophosphate induced neuropsychiatric disorder (COPIND): Results of two postal questionnaire surveys. *J Nutritional & Environmmetal Medicine* 1999;9:123-34.
64. Devanur LD, Kerr JR. Chronic fatigue syndrome. *J Clin Virol* 2006;37:139-50.
65. D'Amelio M, Ricci I, Sacco R, et al. Paraoxonase gene variants are associated with autism in North America, but not in Italy: possible regional specificity in gene-environment interactions. *Mol Psychiatry* 2005;10:1006-16.

66. Kaushik N, Fear D, Richards SC, et al. Gene expression in peripheral blood mononuclear cells from patients with chronic fatigue syndrome. *J Clin Pathol* 2005;58:826-32.
67. Pasca SP, Nemes B, Vlase L, et al. High levels of homocysteine and low serum paraoxonase 1 arylesterase activity in children with autism. *Life Sci* 2006;78:2244-8.
68. Hotopf M, David A, Hull L, Ismail K, Unwin C, Wessely S. Role of vaccinations as risk factors for ill health in veterans of the Gulf war: cross sectional study. *Bmj* 2000;320:1363-7.
69. Schumm WR, Reppert EJ, Jurich AP, et al. Self-reported changes in subjective health and anthrax vaccination as reported by over 900 Persian Gulf War era veterans. *Psychol Rep* 2002;90:639-53.
70. Mahan CM, Kang HK, Dalager NA, Heller JM. Anthrax vaccination and self-reported symptoms, functional status, and medical conditions in the National Health Survey of Gulf War Era Veterans and Their Families. *Ann Epidemiol* 2004;14:81-8.
71. Cherry N, Creed F, Silman A, et al. Health and exposures of United Kingdom Gulf war veterans. Part II: The relation of health to exposure. *Occup Environ Med* 2001;58:299-306.
72. Chalder T, Hotopf M, Unwin C, et al. Prevalence of Gulf war veterans who believe they have Gulf war syndrome: questionnaire study. *Bmj* 2001;323:473-6.
73. Commonwealth Department of Veterans' Affairs. Australian Gulf War Veterans' Health Study 2003. http://www.dv.gov.au/media/publicat/2003/gulfwarhs/html/executive_summaryvhtm 2003.
74. Ortega-Hernandez OD, Shoenfeld Y. Infection, vaccination, and autoantibodies in chronic fatigue syndrome, cause or coincidence? *Ann N Y Acad Sci* 2009;1173:600-9.
75. Nancy AL, Yehuda S. Chronic fatigue syndrome with autoantibodies - The result of an augmented adjuvant effect of hepatitis-B vaccine and silicone implant. *Autoimmun Rev* 2008.
76. Steele J. 2 marines who balked at vaccine deployed. *San Diego Union Tribune* 2003;02-08-2003.
77. Milatovic D, Gupta RC, Aschner M. Anticholinesterase toxicity and oxidative stress. *ScientificWorldJournal* 2006;6:295-310.
78. Li L, Shou Y, Borowitz JL, Isom GE. Reactive oxygen species mediate pyridostigmine-induced neuronal apoptosis: involvement of muscarinic and NMDA receptors. *Toxicol Appl Pharmacol* 2001;177:17-25.
79. Pena-Llopis S, Ferrando MD, Pena JB. Impaired glutathione redox status is associated with decreased survival in two organophosphate-poisoned marine bivalves. *Chemosphere* 2002;47:485-97.
80. Pena-Llopis S, Ferrando MD, Pena JB. Fish tolerance to organophosphate-induced oxidative stress is dependent on the glutathione metabolism and enhanced by N-acetylcysteine. *Aquat Toxicol* 2003;65:337-60.
81. Pena-Llopis S, Ferrando MD, Pena JB. Increased recovery of brain acetylcholinesterase activity in dichlorvos-intoxicated European eels *Anguilla anguilla* by bath treatment with N-acetylcysteine. *Dis Aquat Organ* 2003;55:237-45.
82. Clapp BR, Hingorani AD, Kharbanda RK, et al. Inflammation-induced endothelial dysfunction involves reduced nitric oxide bioavailability and increased oxidant stress. *Cardiovasc Res* 2004;64:172-8.
83. Agrawal A, Chandra D, Kale RK. Radiation induced oxidative stress: II studies in liver as a distant organ of tumor bearing mice. *Mol Cell Biochem* 2001;224:9-17.
84. Leonard SS, Harris GK, Shi X. Metal-induced oxidative stress and signal transduction. *Free Radic Biol Med* 2004;37:1921-42.
85. Piotrowska D, Dlugosz A, Pajak J. Antioxidative properties of coenzyme Q10 and vitamin E in exposure to xylene and gasoline and their mixture with methanol. *Acta Pol Pharm* 2002;59:427-32.
86. Mandavilli BS, Santos JH, Van Houten B. Mitochondrial DNA repair and aging. *Mutat Res* 2002;509:127-51.
87. Genova ML, Pich MM, Bernacchia A, et al. The mitochondrial production of reactive oxygen species in relation to aging and pathology. *Ann N Y Acad Sci* 2004;1011:86-100.
88. Lee HC, Wei YH. Role of Mitochondria in Human Aging. *J Biomed Sci* 1997;4:319-26.
89. Sastre J, Pallardo FV, Vina J. The role of mitochondrial oxidative stress in aging. *Free Radic Biol Med* 2003;35:1-8.

90. Staniek K, Gille L, Kozlov AV, Nohl H. Mitochondrial superoxide radical formation is controlled by electron bifurcation to the high and low potential pathways. *Free Radic Res* 2002;36:381-7.
91. Adam-Vizi V. Production of reactive oxygen species in brain mitochondria: contribution by electron transport chain and non-electron transport chain sources. *Antioxid Redox Signal* 2005;7:1140-9.
92. Wallace DC, Ye JH, Neckelmann SN, Singh G, Webster KA, Greenberg BD. Sequence analysis of cDNAs for the human and bovine ATP synthase beta subunit: mitochondrial DNA genes sustain seventeen times more mutations. *Curr Genet* 1987;12:81-90.
93. Khaidakov M, Heflich RH, Manjanatha MG, Myers MB, Aidoo A. Accumulation of point mutations in mitochondrial DNA of aging mice. *Mutat Res* 2003;526:1-7.
94. Lombes A, Bonilla E, Dimauro S. Mitochondrial encephalomyopathies. *Rev Neurol (Paris)* 1989;145:671-89.
95. Schoeler S, Szibor R, Gellerich FN, et al. Mitochondrial DNA deletions sensitize cells to apoptosis at low heteroplasmy levels. *Biochem Biophys Res Commun* 2005;332:43-9.
96. Wallace DC. The mitochondrial genome in human adaptive radiation and disease: on the road to therapeutics and performance enhancement. *Gene* 2005;354:169-80.
97. Gill MB, Perez-Polo JR. Hypoxia Ischemia-Mediated Cell Death in Neonatal Rat Brain. *Neurochem Res* 2008.
98. Schindowski K, Leutner S, Kressmann S, Eckert A, Muller WE. Age-related increase of oxidative stress-induced apoptosis in mice prevention by Ginkgo biloba extract (EGb761). *J Neural Transm* 2001;108:969-78.
99. Nishio C, Yoshida K, Nishiyama K, Hatanaka H, Yamada M. Involvement of cystatin C in oxidative stress-induced apoptosis of cultured rat CNS neurons. *Brain Res* 2000;873:252-62.
100. Vitale M, Di Matola T, D'Ascoli F, et al. Iodide excess induces apoptosis in thyroid cells through a p53-independent mechanism involving oxidative stress. *Endocrinology* 2000;141:598-605.
101. Lee HC, Wei YH. Mutation and oxidative damage of mitochondrial DNA and defective turnover of mitochondria in human aging. *J Formos Med Assoc* 1997;96:770-8.
102. Wright G, Terada K, Yano M, Sergeev I, Mori M. Oxidative stress inhibits the mitochondrial import of preproteins and leads to their degradation. *Exp Cell Res* 2001;263:107-17.
103. Deth R, Muratore C, Benzecry J, Power-Charnitsky VA, Waly M. How environmental and genetic factors combine to cause autism: A redox/methylation hypothesis. *Neurotoxicology* 2008;29:190-201.
104. Facheris M, Beretta S, Ferrarese C. Peripheral markers of oxidative stress and excitotoxicity in neurodegenerative disorders: tools for diagnosis and therapy? *J Alzheimers Dis* 2004;6:177-84.
105. Jenner P. Oxidative stress in Parkinson's disease. *Ann Neurol* 2003;53 Suppl 3:S26-36; discussion S-8.
106. Chauhan A, Chauhan V. Oxidative stress in autism. *Pathophysiology* 2006;13:171-81.
107. Blaylock RL. A possible central mechanism in autism spectrum disorders, part 2: immunoexcitotoxicity. *Altern Ther Health Med* 2009;15:60-7.
108. De Freitas MR, Chimelli L, Nascimento OJ, Cincinatus D, Marques HA, Nevares MT. [Polyneuropathy caused by trichlorfon: report of a case with electrophysiologic and histopathologic study of the sural nerve]. *Arq Neuropsiquiatr* 1990;48:515-9.
109. Karbowski M, Spodnik JH, Teranishi M, et al. Opposite effects of microtubule-stabilizing and microtubule-destabilizing drugs on biogenesis of mitochondria in mammalian cells. *J Cell Sci* 2001;114:281-91.
110. Knabe W, Kuhn HJ. The role of microtubules and microtubule-organising centres during the migration of mitochondria. *J Anat* 1996;189 (Pt 2):383-91.
111. Chan DC. Mitochondria: dynamic organelles in disease, aging, and development. *Cell* 2006;125:1241-52.
112. Schuman E, Chan D. Fueling synapses. *Cell* 2004;119:738-40.
113. Tissandie E, Gueguen Y, Lobaccaro JM, et al. In vivo effects of chronic contamination with depleted uranium on vitamin D3 metabolism in rat. *Biochim Biophys Acta* 2007;1770:266-72.

114. Crivello JF. Oxidative stress limits vitamin D metabolism by bovine proximal tubule cells in vitro. *Arch Biochem Biophys* 1988;262:471-80.
115. Adorini L. Immunomodulatory effects of vitamin D receptor ligands in autoimmune diseases. *Int Immunopharmacol* 2002;2:1017-28.
116. Cantorna MT, Mahon BD. Mounting evidence for vitamin D as an environmental factor affecting autoimmune disease prevalence. *Exp Biol Med (Maywood)* 2004;229:1136-42.
117. Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* 2004;80:1678S-88S.
118. Holick MF. Vitamin D: important for prevention of osteoporosis, cardiovascular heart disease, type 1 diabetes, autoimmune diseases, and some cancers. *South Med J* 2005;98:1024-7.
119. Meehan TF, DeLuca HF. The vitamin D receptor is necessary for 1alpha,25-dihydroxyvitamin D(3) to suppress experimental autoimmune encephalomyelitis in mice. *Arch Biochem Biophys* 2002;408:200-4.
120. Vojdani A, Thrasher JD. Cellular and humoral immune abnormalities in Gulf War veterans. *Environ Health Perspect* 2004;112:840-6.
121. Asa PB, Cao Y, Garry RF. Antibodies to squalene in Gulf War syndrome. *Exp Mol Pathol* 2000;68:55-64.
122. Exley C, Swarbrick L, Gherardi RK, Authier FJ. A role for the body burden of aluminium in vaccine-associated macrophagic myofasciitis and chronic fatigue syndrome. *Med Hypotheses* 2009;72:135-9.
123. McDonald T. MOD Anthrax Vaccine Contains Squalene. "Tonight with Trevor McDonald", March 17, 2003, ITV1, 8PM 2003 Mar 17, 2003.
124. Rodriguez PM. Squalene presence confirmed by FDA. *Insight* 2000;<http://www.insightmag.com/news/2000/10/30/TheLastWord/Squalene.Presence.Confirmed.By.Fda-213345.shtml>.
125. Bonnefont-Rousselot D, Chantalat-Auger C, Teixeira A, Jaudon MC, Pelletier S, Cherin P. Blood oxidative stress status in patients with macrophagic myofasciitis. *Biomed Pharmacother* 2004;58:516-9.
126. Gherardi RK. [Lessons from macrophagic myofasciitis: towards definition of a vaccine adjuvant-related syndrome]. *Rev Neurol (Paris)* 2003;159:162-4.
127. Gherardi RK, Coquet M, Cherin P, et al. Macrophagic myofasciitis: an emerging entity. Group d'Etudes et Recherche sur les Maladies Musculaires Acquises et Dysimmunitaires (GERMMAD) de l'Association Francaise contre les Myopathies (AFM). *Lancet* 1998;352:347-52.
128. Shoenfeld Y, Agmon-Levin N. 'ASIA' - Autoimmune/inflammatory syndrome induced by adjuvants. *J Autoimmun* 2010.
129. Neaton JD, Wentworth DN, Kuller LH. Relationship of serum cholesterol and blood pressure with risk of death from AIDS [Abstract]. GET FULL REFERENCE FROM MHC 199?:6-7.
130. Jacobs DRJ, Sidney S, Feingold K, Iribarren C. Inverse association between blood total cholesterol and HIV infection, AIDS, and AIDS death: Cause or effect? [Abstract]. GET JOURNAL; REFERENCE NOT STATED ON PAGE 199?:5-6.
131. Villamor E. A potential role for vitamin D on HIV infection? *Nutr Rev* 2006;64:226-33.
132. Ginde AA, Mansbach JM, Camargo CA, Jr. Vitamin D, respiratory infections, and asthma. *Curr Allergy Asthma Rep* 2009;9:81-7.
133. Walker VP, Modlin RL. The Vitamin D Connection to Pediatric Infections and Immune Function. *Pediatr Res* 2009.
134. Li JH, Chen DM, Li Z, et al. [Study on association between vitamin D receptor gene polymorphisms and the outcomes of HBV infection]. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 2006;23:402-5.
135. Nakamura K, Matsunaga K. Susceptibility of natural killer (NK) cells to reactive oxygen species (ROS) and their restoration by the mimics of superoxide dismutase (SOD). *Cancer Biother Radiopharm* 1998;13:275-90.

136. Geier MR, Geier DA. A case-series of adverse events, positive re-challenge of symptoms, and events in identical twins following hepatitis B vaccination: analysis of the Vaccine Adverse Event Reporting System (VAERS) database and literature review. *Clin Exp Rheumatol* 2004;22:749-55.
137. Asa PB, Wilson RB, Garry RF. Antibodies to squalene in recipients of anthrax vaccine. *Exp Mol Pathol* 2002;73:19-27.
138. Kerrison JB, Lounsbury D, Thirkill CE, Lane RG, Schatz MP, Engler RM. Optic neuritis after anthrax vaccination. *Ophthalmology* 2002;109:99-104.
139. Lane RG, et al. Optic neuritis following anthrax vaccination. *Iovs* 2001;42:S326.
140. Hakariya Y, Kuratsune H. [Chronic fatigue syndrome: biochemical examination of blood]. *Nippon Rinsho* 2007;65:1071-6.
141. Singh VK, Rivas WH. Detection of antinuclear and antilaminin antibodies in autistic children who received thimerosal-containing vaccines. *J Biomed Sci* 2004;11:607-10.
142. Singh VK. Thimerosal is unrelated to autoimmune autism. *Pediatr Allergy Immunol* 2007;18:89.
143. Valicenti-McDermott MD, McVicar K, Cohen HJ, Wershil BK, Shinnar S. Gastrointestinal symptoms in children with an autism spectrum disorder and language regression. *Pediatr Neurol* 2008;39:392-8.
144. Castilho RF, Ward MW, Nicholls DG. Oxidative stress, mitochondrial function, and acute glutamate excitotoxicity in cultured cerebellar granule cells. *J Neurochem* 1999;72:1394-401.
145. Singh P, Mann KA, Mangat HK, Kaur G. Prolonged glutamate excitotoxicity: effects on mitochondrial antioxidants and antioxidant enzymes. *Mol Cell Biochem* 2003;243:139-45.
146. Mackness B, Mackness MI, Arrol S, Turkie W, Durrington PN. Effect of the molecular polymorphisms of human paraoxonase (PON1) on the rate of hydrolysis of paraoxon. *Br J Pharmacol* 1997;122:265-8.
147. Schnakenberg E, Fabig KR, Stanulla M, et al. A cross-sectional study of self-reported chemical-related sensitivity is associated with gene variants of drug-metabolizing enzymes. *Environ Health* 2007;6:6.
148. James SJ, Melnyk S, Jernigan S, et al. Metabolic endophenotype and related genotypes are associated with oxidative stress in children with autism. *Am J Med Genet B Neuropsychiatr Genet* 2006;141B:947-56.
149. James SJ, Cutler P, Melnyk S, et al. Metabolic biomarkers of increased oxidative stress and impaired methylation capacity in children with autism. *Am J Clin Nutr* 2004;80:1611-7.
150. Sogut S, Zoroglu SS, Ozyurt H, et al. Changes in nitric oxide levels and antioxidant enzyme activities may have a role in the pathophysiological mechanisms involved in autism. *Clin Chim Acta* 2003;331:111-7.
151. Hotopf M, Mackness MI, Nikolaou V, et al. Paraoxonase in Persian Gulf War veterans. *J Occup Environ Med* 2003;45:668-75.
152. Lee HA, Bale AJ, Gabriel R. Results of investigations on Gulf War veterans. *Clin Med* 2005;5:166-72.
153. Ismail K, Kent K, Sherwood R, et al. Chronic fatigue syndrome and related disorders in UK veterans of the Gulf War 1990-1991: results from a two-phase cohort study. *Psychol Med* 2007;1-9.
154. Whistler T, Fletcher MA, Lonergan W, et al. Impaired immune function in Gulf War Illness. *BMC Med Genomics* 2009;2:12.
155. Haley RW, Vongpatanasin W, Wolfe GI, et al. Blunted circadian variation in autonomic regulation of sinus node function in veterans with Gulf War syndrome. *Am J Med* 2004;117:469-78.
156. Stein PK, Domitrovich PP, Ambrose K, et al. Sex effects on heart rate variability in fibromyalgia and Gulf War illness. *Arthritis Rheum* 2004;51:700-8.
157. Hannan KL, Berg DE, Baumzweiger W, et al. Activation of the coagulation system in Gulf War Illness: a potential pathophysiologic link with chronic fatigue syndrome. A laboratory approach to diagnosis. *Blood Coagul Fibrinolysis* 2000;11:673-8.

158. Bach R. Gulf War associated chronic coagulopathies update. In: Meeting of the Research Advisory Committee on Gulf War Veterans Illnesses; 2009 June 29; Boston University School of Public Health; 2009.
159. Hopps E, Noto D, Caimi G, Averna MR. A novel component of the metabolic syndrome: the oxidative stress. *Nutr Metab Cardiovasc Dis* 2010;20:72-7.
160. Thomas HV, Stimpson NJ, Weightman AL, Dunstan F, Lewis G. Systematic review of multi-symptom conditions in Gulf War veterans. *Psychol Med* 2006;36:735-47.
161. Bourdette DN, McCauley LA, Barkhuizen A, et al. Symptom factor analysis, clinical findings, and functional status in a population-based case control study of Gulf War unexplained illness. *J Occup Environ Med* 2001;43:1026-40.
162. Baraniuk JN, Casado B, Maibach H, Clauw DJ, Pannell LK, Hess SS. A Chronic Fatigue Syndrome - related proteome in human cerebrospinal fluid. *BMC Neurol* 2005;5:22.
163. Ciccone DS, Weissman L, Natelson BH. Chronic fatigue syndrome in male Gulf war veterans and civilians: a further test of the single syndrome hypothesis. *J Health Psychol* 2008;13:529-36.
164. Eisen SA, Kang HK, Murphy FM, et al. Gulf War veterans' health: medical evaluation of a U.S. cohort. *Ann Intern Med* 2005;142:881-90.
165. Jammes Y, Steinberg JG, Mambrini O, Bregeon F, Delliaux S. Chronic fatigue syndrome: assessment of increased oxidative stress and altered muscle excitability in response to incremental exercise. *J Intern Med* 2005;257:299-310.
166. Manuel y Keenoy B, Moorkens G, Vertommen J, De Leeuw I. Antioxidant status and lipoprotein peroxidation in chronic fatigue syndrome. *Life Sci* 2001;68:2037-49.
167. Kennedy G, Spence VA, McLaren M, Hill A, Underwood C, Belch JJ. Oxidative stress levels are raised in chronic fatigue syndrome and are associated with clinical symptoms. *Free Radic Biol Med* 2005;39:584-9.
168. Plioplys AV, Plioplys S. Serum levels of carnitine in chronic fatigue syndrome: clinical correlates. *Neuropsychobiology* 1995;32:132-8.
169. Kuratsune H, Yamaguti K, Takahashi M, Misaki H, Tagawa S, Kitani T. Acylcarnitine deficiency in chronic fatigue syndrome. *Clin Infect Dis* 1994;18 Suppl 1:S62-7.
170. Lane RJ, Barrett MC, Taylor DJ, Kemp GJ, Lodi R. Heterogeneity in chronic fatigue syndrome: evidence from magnetic resonance spectroscopy of muscle. *Neuromuscul Disord* 1998;8:204-9.
171. Chaudhuri A, Behan PO. In vivo magnetic resonance spectroscopy in chronic fatigue syndrome. *Prostaglandins Leukot Essent Fatty Acids* 2004;71:181-3.
172. Myhill S, Booth NE, McLaren-Howard J. Chronic fatigue syndrome and mitochondrial dysfunction. *Int J Clin Exp Med* 2009;2:1-16.
173. Whistler T, Unger ER, Nisenbaum R, Vernon SD. Integration of gene expression, clinical, and epidemiologic data to characterize Chronic Fatigue Syndrome. *J Transl Med* 2003;1:10.
174. Fang H, Xie Q, Boneva R, Fostel J, Perkins R, Tong W. Gene expression profile exploration of a large dataset on chronic fatigue syndrome. *Pharmacogenomics* 2006;7:429-40.
175. Benito-Leon J, Berbel A, Porta-Estessam J, Martinez A, Arenas J. [Fibromyalgia in right half of the body as the onset of mitochondrial cytopathy. Letter]. *Rev Neurol* 1996;24:1303-4.
176. Villanova M, Selvi E, Malandrini A, et al. Mitochondrial myopathy mimicking fibromyalgia syndrome. *Muscle Nerve* 1999;22:289-91.
177. Bengtsson A, Henriksson KG. The muscle in fibromyalgia--a review of Swedish studies. *J Rheumatol Suppl* 1989;19:144-9.
178. Dufault R, Schnoll R, Lukiw WJ, et al. Mercury exposure, nutritional deficiencies and metabolic disruptions may affect learning in children. *Behav Brain Funct* 2009;5:44.
179. Geier DA, Kern JK, Garver CR, Adams JB, Audhya T, Geier MR. A Prospective Study of Transsulfuration Biomarkers in Autistic Disorders. *Neurochem Res* 2008.
180. Geier DA, Kern JK, Garver CR, et al. Biomarkers of environmental toxicity and susceptibility in autism. *J Neurol Sci* 2008.

181. Poling JS, Frye RE, Shoffner J, Zimmerman AW. Developmental regression and mitochondrial dysfunction in a child with autism. *J Child Neurol* 2006;21:170-2.
182. Kern JK, Jones AM. Evidence of toxicity, oxidative stress, and neuronal insult in autism. *J Toxicol Environ Health B Crit Rev* 2006;9:485-99.
183. Loevinger BL, Muller D, Alonso C, Coe CL. Metabolic syndrome in women with chronic pain. *Metabolism* 2007;56:87-93.
184. Bell IR, Warg-Damiani L, Baldwin CM, Walsh ME, Schwartz GE. Self-reported chemical sensitivity and wartime chemical exposures in Gulf War veterans with and without decreased global health ratings. *Mil Med* 1998;163:725-32.
185. Reid S, Hotopf M, Hull L, Ismail K, Unwin C, Wessely S. Multiple chemical sensitivity and chronic fatigue syndrome in British Gulf War veterans. *Am J Epidemiol* 2001;153:604-9.
186. Aaron LA, Buchwald D. Chronic diffuse musculoskeletal pain, fibromyalgia and co-morbid unexplained clinical conditions. *Best Pract Res Clin Rheumatol* 2003;17:563-74.
187. Tafazoli S, O'Brien PJ. Peroxidases: a role in the metabolism and side effects of drugs. *Drug Discov Today* 2005;10:617-25.
188. Tafazoli S, Spehar DD, O'Brien PJ. Oxidative stress mediated idiosyncratic drug toxicity. *Drug Metab Rev* 2005;37:311-25.
189. Perez-Gomez C, Segura JM, Blanca M, Asenjo M, Mates JM. Antioxidant activity levels and oxidative stress as blood markers of allergic response to drugs. *Biochem Cell Biol* 2000;78:691-8.
190. Kass GE. Mitochondrial involvement in drug-induced hepatic injury. *Chem Biol Interact* 2006;163:145-59.
191. McKeown-Eyssen G, Baines C, Cole DE, et al. Case-control study of genotypes in multiple chemical sensitivity: CYP2D6, NAT1, NAT2, PON1, PON2 and MTHFR. *Int J Epidemiol* 2004;33:971-8.
192. Haley RW, Kurt TL. Self-reported exposure to neurotoxic chemical combinations in the Gulf War. A cross-sectional epidemiologic study. *Jama* 1997;277:231-7.
193. Schumm WR, Reppert EJ, Jurich AP, et al. Pyridostigmine bromide and the long-term subjective health status of a sample of over 700 male Reserve Component Gulf War era veterans. *Psychol Rep* 2002;90:707-21.
194. Liu X, Hubbard JA, Fabes RA, Adam JB. Sleep disturbances and correlates of children with autism spectrum disorders. *Child Psychiatry Hum Dev* 2006;37:179-91.
195. de Magistris L, Familiari V, Pascotto A, et al. Alterations of the intestinal barrier in patients with autism spectrum disorders and in their first-degree relatives. *J Pediatr Gastroenterol Nutr* 2010;51:418-24.
196. Smith RA, Farnworth H, Wright B, Allgar V. Are there more bowel symptoms in children with autism compared to normal children and children with other developmental and neurological disorders?: A case control study. *Autism* 2009;13:343-55.
197. Jyonouchi H, Geng L, Ruby A, Reddy C, Zimmerman-Bier B. Evaluation of an association between gastrointestinal symptoms and cytokine production against common dietary proteins in children with autism spectrum disorders. *J Pediatr* 2005;146:605-10.
198. Jill James S, Melnyk S, Jernigan S, Hubanks A, Rose S, Gaylor DW. Abnormal Transmethylation/transsulfuration Metabolism and DNA Hypomethylation Among Parents of Children with Autism. *J Autism Dev Disord* 2008.
199. Correia C, Coutinho AM, Diogo L, et al. Brief report: High frequency of biochemical markers for mitochondrial dysfunction in autism: no association with the mitochondrial aspartate/glutamate carrier SLC25A12 gene. *J Autism Dev Disord* 2006;36:1137-40.
200. Oliveira G, Diogo L, Grazina M, et al. Mitochondrial dysfunction in autism spectrum disorders: a population-based study. *Dev Med Child Neurol* 2005;47:185-9.
201. Ming X, Stein TP, Brimacombe M, Johnson WG, Lambert GH, Wagner GC. Increased excretion of a lipid peroxidation biomarker in autism. *Prostaglandins Leukot Essent Fatty Acids* 2005;73:379-84.

202. Zoroglu SS, Armutcu F, Ozen S, et al. Increased oxidative stress and altered activities of erythrocyte free radical scavenging enzymes in autism. *Eur Arch Psychiatry Clin Neurosci* 2004;254:143-7.
203. Ramoz N, Reichert JG, Smith CJ, et al. Linkage and association of the mitochondrial aspartate/glutamate carrier SLC25A12 gene with autism. *Am J Psychiatry* 2004;161:662-9.
204. Filipek PA, Juranek J, Nguyen MT, Cummings C, Gargus JJ. Relative carnitine deficiency in autism. *J Autism Dev Disord* 2004;34:615-23.
205. Chauhan A, Chauhan V, Brown WT, Cohen I. Oxidative stress in autism: increased lipid peroxidation and reduced serum levels of ceruloplasmin and transferrin--the antioxidant proteins. *Life Sci* 2004;75:2539-49.
206. Blaxill MF, Redwood L, Bernard S. Thimerosal and autism? A plausible hypothesis that should not be dismissed. *Med Hypotheses* 2004;62:788-94.
207. Lombard J. Autism: a mitochondrial disorder? *Med Hypotheses* 1998;50:497-500.
208. Patarca R. Cytokines and chronic fatigue syndrome. *Ann N Y Acad Sci* 2001;933:185-200.
209. Brenu EW, van Driel ML, Staines DR, et al. Immunological abnormalities as potential biomarkers in Chronic Fatigue Syndrome/Myalgic Encephalomyelitis. *J Transl Med* 2011;9:81.
210. Berg D, Berg LH, Couvaras J, Harrison H. Chronic fatigue syndrome and/or fibromyalgia as a variation of antiphospholipid antibody syndrome: an explanatory model and approach to laboratory diagnosis. *Blood Coagul Fibrinolysis* 1999;10:435-8.
211. Fix Author List, Menon DU. Biochemical screening for mitochondrial disease in patients with autism spectrum disorder. In: *AMFAR*; 2011; San Diego; 2011.
212. Geier MR, Geier DA. Gastrointestinal adverse reactions following anthrax vaccination: an analysis of the Vaccine Adverse Events Reporting System (VAERS) database. *Hepatogastroenterology* 2004;51:762-7.
213. De Marchi U, Mancon M, Battaglia V, Ceccon S, Cardellini P, Toninello A. Influence of reactive oxygen species production by monoamine oxidase activity on aluminum-induced mitochondrial permeability transition. *Cell Mol Life Sci* 2004;61:2664-71.
214. Gomez M, Esparza JL, Nogues MR, Giralt M, Cabre M, Domingo JL. Pro-oxidant activity of aluminum in the rat hippocampus: gene expression of antioxidant enzymes after melatonin administration. *Free Radic Biol Med* 2005;38:104-11.
215. Jyoti A, Sharma D. Neuroprotective role of Bacopa monniera extract against aluminium-induced oxidative stress in the hippocampus of rat brain. *Neurotoxicology* 2006;27:451-7.
216. Katyal R, Desigan B, Sodhi CP, Ojha S. Oral aluminum administration and oxidative injury. *Biol Trace Elem Res* 1997;57:125-30.
217. Lankoff A, Banasik A, Duma A, et al. A comet assay study reveals that aluminium induces DNA damage and inhibits the repair of radiation-induced lesions in human peripheral blood lymphocytes. *Toxicol Lett* 2006;161:27-36.
218. Mailloux R, Lemire J, Appanna V. Aluminum-induced mitochondrial dysfunction leads to lipid accumulation in human hepatocytes: a link to obesity. *Cell Physiol Biochem* 2007;20:627-38.
219. Murakami K, Yoshino M. Aluminum decreases the glutathione regeneration by the inhibition of NADP-isocitrate dehydrogenase in mitochondria. *J Cell Biochem* 2004;93:1267-71.
220. Nehru B, Anand P. Oxidative damage following chronic aluminium exposure in adult and pup rat brains. *J Trace Elem Med Biol* 2005;19:203-8.
221. Sargazi M, Shenkin A, Roberts NB. Aluminium-induced injury to kidney proximal tubular cells: Effects on markers of oxidative damage. *J Trace Elem Med Biol* 2006;19:267-73.
222. Orihuela D, Meichtry V, Pregi N, Pizarro M. Short-term oral exposure to aluminium decreases glutathione intestinal levels and changes enzyme activities involved in its metabolism. *J Inorg Biochem* 2005;99:1871-8.
223. Messer RL, Lockwood PE, Tseng WY, et al. Mercury (II) alters mitochondrial activity of monocytes at sublethal doses via oxidative stress mechanisms. *J Biomed Mater Res B Appl Biomater* 2005;75:257-63.

224. Chen C, Qu L, Li B, et al. Increased oxidative DNA damage, as assessed by urinary 8-hydroxy-2'-deoxyguanosine concentrations, and serum redox status in persons exposed to mercury. *Clin Chem* 2005;51:759-67.
225. Ueha-Ishibashi T, Oyama Y, Nakao H, et al. Effect of thimerosal, a preservative in vaccines, on intracellular Ca²⁺ concentration of rat cerebellar neurons. *Toxicology* 2004;195:77-84.
226. Makani S, Gollapudi S, Yel L, Chiplunkar S, Gupta S. Biochemical and molecular basis of thimerosal-induced apoptosis in T cells: a major role of mitochondrial pathway. *Genes Immun* 2002;3:270-8.
227. El-Demerdash FM. Effects of selenium and mercury on the enzymatic activities and lipid peroxidation in brain, liver, and blood of rats. *J Environ Sci Health B* 2001;36:489-99.
228. Pizzichini M, Fonzi M, Sugherini L, et al. Release of mercury from dental amalgam and its influence on salivary antioxidant activity. *Bull Group Int Rech Sci Stomatol Odontol* 2000;42:94-100.
229. Chung CP, Titova D, Oeser A, et al. Oxidative stress in fibromyalgia and its relationship to symptoms. *Clin Rheumatol* 2008.
230. Hooper RG, Thomas AR, Kearl RA. Mitochondrial enzyme deficiency causing exercise limitation in normal-appearing adults. *Chest* 1995;107:317-22.
231. Smeitink JA. Mitochondrial disorders: clinical presentation and diagnostic dilemmas. *J Inherit Metab Dis* 2003;26:199-207.
232. Fadic R, Johns DR. Clinical spectrum of mitochondrial diseases. *Semin Neurol* 1996;16:11-20.
233. Zeviani M, Antozzi C. Mitochondrial disorders. *Mol Hum Reprod* 1997;3:133-48.
234. Jeyakumar A, Williamson ME, Brickman TM, Krakovitz P, Parikh S. Otolaryngologic manifestations of mitochondrial cytopathies. *Am J Otolaryngol* 2009;30:162-5.
235. Banerjee BD, Seth V, Bhattacharya A, Pasha ST, Chakraborty AK. Biochemical effects of some pesticides on lipid peroxidation and free-radical scavengers. *Toxicol Lett* 1999;107:33-47.
236. Whitfield JB. Gamma glutamyl transferase. *Crit Rev Clin Lab Sci* 2001;38:263-355.
237. Bajnok L, Seres I, Varga Z, et al. Relationship of serum resistin level to traits of metabolic syndrome and serum paraoxonase 1 activity in a population with a broad range of body mass index. *Exp Clin Endocrinol Diabetes* 2008;116:592-9.
238. Serhatlioglu S, Gursu MF, Gulcu F, Canatan H, Godekmerdan A. Levels of paraoxonase and arylesterase activities and malondialdehyde in workers exposed to ionizing radiation. *Cell Biochem Funct* 2003;21:371-5.
239. Viora M, Quaranta MG, Straface E, Vari R, Masella R, Malorni W. Redox imbalance and immune functions: opposite effects of oxidized low-density lipoproteins and N-acetylcysteine. *Immunology* 2001;104:431-8.
240. Droge W, Breitkreutz R. Glutathione and immune function. *Proc Nutr Soc* 2000;59:595-600.
241. Wood SM, Beckham C, Yosioka A, Darban H, Watson RR. beta-Carotene and selenium supplementation enhances immune response in aged humans. *Integr Med* 2000;2:85-92.
242. Manzella D, Barbieri M, Ragno E, Paolisso G. Chronic administration of pharmacologic doses of vitamin E improves the cardiac autonomic nervous system in patients with type 2 diabetes. *Am J Clin Nutr* 2001;73:1052-7.
243. Schwartz J, Park SK, O'Neill MS, et al. Glutathione-S-transferase M1, obesity, statins, and autonomic effects of particles: gene-by-drug-by-environment interaction. *Am J Respir Crit Care Med* 2005;172:1529-33.
244. Davis SD, Kator SF, Wonnott JA, Pappas BL, Sall JL. Neurally mediated hypotension in fatigued Gulf War veterans: a preliminary report. *Am J Med Sci* 2000;319:89-95.
245. Sastre A. Research Presentation. In: Meeting of the Research Advisory Committee on Gulf War Veterans' Illnesses (Public meeting); 2003 June 16
Department of Veterans Affairs, Washington, D.C., Room 230; 2003.
246. Majamaa-Voltti K, Majamaa K, Peuhkurinen K, Makikallio TH, Huikuri HV. Cardiovascular autonomic regulation in patients with 3243A > G mitochondrial DNA mutation. *Ann Med* 2004;36:225-31.

247. Momiyama Y, Suzuki Y, Ohtomo M, et al. Cardiac autonomic nervous dysfunction in diabetic patients with a mitochondrial DNA mutation: assessment by heart rate variability. *Diabetes Care* 2002;25:2308-13.
248. Mezzano D, Pais EO, Aranda E, et al. Inflammation, not hyperhomocysteinemia, is related to oxidative stress and hemostatic and endothelial dysfunction in uremia. *Kidney Int* 2001;60:1844-50.
249. Ceriello A. Coagulation activation in diabetes mellitus: the role of hyperglycaemia and therapeutic prospects. *Diabetologia* 1993;36:1119-25.
250. Pawlak K, Naumnik B, Brzosko S, Pawlak D, Mysliwiec M. Oxidative stress - a link between endothelial injury, coagulation activation, and atherosclerosis in haemodialysis patients. *Am J Nephrol* 2004;24:154-61.
251. Ceriello A, Bortolotti N, Motz E, et al. Red wine protects diabetic patients from meal-induced oxidative stress and thrombosis activation: a pleasant approach to the prevention of cardiovascular disease in diabetes. *Eur J Clin Invest* 2001;31:322-8.
252. Sastre J, Pallardo FV, Vina J. Mitochondrial oxidative stress plays a key role in aging and apoptosis. *IUBMB Life* 2000;49:427-35.
253. Reutelingsperger CP, van Heerde WL. Annexin V, the regulator of phosphatidylserine-catalyzed inflammation and coagulation during apoptosis. *Cell Mol Life Sci* 1997;53:527-32.
254. Kalousova M, Fialova L, Skrha J, et al. Oxidative stress, inflammation and autoimmune reaction in type 1 and type 2 diabetes mellitus. *Prague Med Rep* 2004;105:21-8.
255. Resch U, Tatzber F, Budinsky A, Sinzinger H. Reduction of oxidative stress and modulation of autoantibodies against modified low-density lipoprotein after rosuvastatin therapy. *Br J Clin Pharmacol* 2006;61:262-74.
256. Traverso N, Patriarca S, Balbis E, et al. Anti malondialdehyde-adduct immunological response as a possible marker of successful aging. *Exp Gerontol* 2003;38:1129-35.
257. Gearhart DA, Sickles DW, Buccafusco JJ, Prendergast MA, Terry AV, Jr. Chlorpyrifos, chlorpyrifos-oxon, and diisopropylfluorophosphate inhibit kinesin-dependent microtubule motility. *Toxicol Appl Pharmacol* 2007;218:20-9.
258. Prendergast MA, Self RL, Smith KJ, et al. Microtubule-associated targets in chlorpyrifos oxon hippocampal neurotoxicity. *Neuroscience* 2007;146:330-9.
259. Golomb BA. A Review of the Scientific Literature as it Pertains to Gulf War Illnesses, Vol 3: Immunizations. Santa Monica: RAND; in press.
260. Pizzichini M, Fonzi M, Sugherini L, et al. Release of mercury from dental amalgam and its influence on salivary antioxidant activity. *Sci Total Environ* 2002;284:19-25.
261. Sharma P, Mishra KP. Aluminum-induced maternal and developmental toxicity and oxidative stress in rat brain: Response to combined administration of Tiron and glutathione. *Reprod Toxicol* 2006;21:313-21.
262. Ueha-Ishibashi T, Tatsuishi T, Iwase K, et al. Property of thimerosal-induced decrease in cellular content of glutathione in rat thymocytes: a flow cytometric study with 5-chloromethylfluorescein diacetate. *Toxicol In Vitro* 2004;18:563-9.
263. Bagchi D, Bagchi M, Hassoun E, Moser J, Stohs SJ. Effects of carbon tetrachloride, menadione, and paraquat on the urinary excretion of malondialdehyde, formaldehyde, acetaldehyde, and acetone in rats. *J Biochem Toxicol* 1993;8:101-6.
264. Bagchi D, Bagchi M, Hassoun EA, Stohs SJ. In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicology* 1995;104:129-40.
265. Sutcu R, Altuntas I, Yildirim B, Karahan N, Demirin H, Delibas N. The effects of subchronic methidathion toxicity on rat liver: Role of antioxidant vitamins C and E. *Cell Biol Toxicol* 2006.
266. Jean JC, Liu Y, Brown LA, Marc RE, Klings E, Joyce-Brady M. Gamma-glutamyl transferase deficiency results in lung oxidant stress in normoxia. *Am J Physiol Lung Cell Mol Physiol* 2002;283:L766-76.

267. Bo S, Gambino R, Durazzo M, et al. Associations between gamma-glutamyl transferase, metabolic abnormalities and inflammation in healthy subjects from a population-based cohort: a possible implication for oxidative stress. *World J Gastroenterol* 2005;11:7109-17.
268. Altuntas I, Delibas N, Demirci M, Kilinc I, Tamer N. The effects of methidathion on lipid peroxidation and some liver enzymes: role of vitamins E and C. *Arch Toxicol* 2002;76:470-3.
269. Mackness B, Durrington PN, Mackness MI. Human serum paraoxonase. *Gen Pharmacol* 1998;31:329-36.
270. Pawlak K, Pawlak D, Mysliwiec M. Extrinsic coagulation pathway activation and metalloproteinase-2/TIMPs system are related to oxidative stress and atherosclerosis in hemodialysis patients. *Thromb Haemost* 2004;92:646-53.
271. Wolvetang EJ, Johnson KL, Krauer K, Ralph SJ, Linnane AW. Mitochondrial respiratory chain inhibitors induce apoptosis. *FEBS Lett* 1994;339:40-4.
272. Xie L, Zhu X, Hu Y, et al. Mitochondrial DNA oxidative damage triggers mitochondrial dysfunction and apoptosis in high glucose-induced human retinal vascular endothelial cells. *Invest Ophthalmol Vis Sci* 2008.
273. Self-reported illness and health status among Gulf War veterans. A population-based study. The Iowa Persian Gulf Study Group. *Jama* 1997;277:238-45.
274. Land JM, Kemp GJ, Taylor DJ, Standing SJ, Radda GK, Rajagopalan B. Oral phosphate supplements reverse skeletal muscle abnormalities in a case of chronic fatigue with idiopathic renal hypophosphatemia. *Neuromuscul Disord* 1993;3:223-5.
275. Barnes PR, Taylor DJ, Kemp GJ, Radda GK. Skeletal muscle bioenergetics in the chronic fatigue syndrome. *J Neurol Neurosurg Psychiatry* 1993;56:679-83.
276. Zhang C, Baumer A, Mackay IR, Linnane AW, Nagley P. Unusual pattern of mitochondrial DNA deletions in skeletal muscle of an adult human with chronic fatigue syndrome. *Hum Mol Genet* 1995;4:751-4.
277. Ozgocmen S, Ozyurt H, Sogut S, Akyol O. Current concepts in the pathophysiology of fibromyalgia: the potential role of oxidative stress and nitric oxide. *Rheumatol Int* 2006;26:585-97.
278. Pongratz DE, Spath M. Morphologic aspects of fibromyalgia. *Z Rheumatol* 1998;57 Suppl 2:47-51.
279. Maes M. Inflammatory and oxidative and nitrosative stress pathways underpinning chronic fatigue, somatization and psychosomatic symptoms. *Curr Opin Psychiatry* 2009;22:75-83.
280. Bradstreet JJ, Smith S, Baral M, Rossignol DA. Biomarker-guided interventions of clinically relevant conditions associated with autism spectrum disorders and attention deficit hyperactivity disorder. *Altern Med Rev* 2010;15:15-32.
281. Giulivi C, Zhang Y-F, Omanska-Klusck A, et al. Mitochondrial dysfunction in autism. *Jama* 2010;304:2389-96.
282. McCauley L, Lasarev M, Sticker D, Rischitelli D, Spencer P. Illness experience of Gulf War veterans possibly exposed to chemical warfare agents. *Am J Prev Med* 2002;23:200.
283. Kang HK, Mahan CM, Lee KY, Magee CA, Murphy FM. Illnesses among United States veterans of the Gulf War: a population-based survey of 30,000 veterans. *J Occup Environ Med* 2000;42:491-501.
284. Johnstone M. Mitochondrial calcium overload, a cause of essential hypertension. *Lancet* 1977;1:650-1.
285. Sun F, Cui J, Gavras H, Schwartz F. A novel class of tests for the detection of mitochondrial DNA-mutation involvement in diseases. *Am J Hum Genet* 2003;72:1515-26.
286. Watson B, Jr., Khan MA, Desmond RA, Bergman S. Mitochondrial DNA mutations in black Americans with hypertension-associated end-stage renal disease. *Am J Kidney Dis* 2001;38:529-36.
287. Lee HK, Song JH, Shin CS, et al. Decreased mitochondrial DNA content in peripheral blood precedes the development of non-insulin-dependent diabetes mellitus. *Diabetes Res Clin Pract* 1998;42:161-7.
288. Arruda WO, Torres LF, Lombes A, et al. Mitochondrial myopathy and myoclonic epilepsy. *Arq Neuropsiquiatr* 1990;48:32-43.

289. Wilson FH, Hariri A, Farhi A, et al. A Cluster of Metabolic Defects Caused by Mutation in a Mitochondrial tRNA. *Science* 2004.
290. Esposito K, Ciotola M, Schisano B, et al. Oxidative stress in the metabolic syndrome. *J Endocrinol Invest* 2006;29:791-5.
291. Fujita K, Nishizawa H, Funahashi T, Shimomura I, Shimabukuro M. Systemic oxidative stress is associated with visceral fat accumulation and the metabolic syndrome. *Circ J* 2006;70:1437-42.
292. Weng SW, Liou CW, Lin TK, et al. Association of mitochondrial deoxyribonucleic acid 16189 variant (T->C transition) with metabolic syndrome in Chinese adults. *J Clin Endocrinol Metab* 2005;90:5037-40.
293. Nicolson GL. Metabolic syndrome and mitochondrial function: molecular replacement and antioxidant supplements to prevent membrane peroxidation and restore mitochondrial function. *J Cell Biochem* 2007;100:1352-69.
294. Coffman CJ, Horner RD, Grambow SC, Lindquist J. Estimating the occurrence of amyotrophic lateral sclerosis among Gulf War (1990-1991) veterans using capture-recapture methods. *Neuroepidemiology* 2005;24:141-50.
295. Haley RW. Excess incidence of ALS in young Gulf War veterans. *Neurology* 2003;61:750-6.
296. Horner RD, Kamins KG, Feussner JR, et al. Occurrence of amyotrophic lateral sclerosis among Gulf War veterans. *Neurology* 2003;61:742-9.
297. Bowling AC, Schulz JB, Brown RH, Jr., Beal MF. Superoxide dismutase activity, oxidative damage, and mitochondrial energy metabolism in familial and sporadic amyotrophic lateral sclerosis. *J Neurochem* 1993;61:2322-5.
298. Ihara Y, Nobukuni K, Takata H, Hayabara T. Oxidative stress and metal content in blood and cerebrospinal fluid of amyotrophic lateral sclerosis patients with and without a Cu, Zn-superoxide dismutase mutation. *Neurol Res* 2005;27:105-8.
299. Ilieva EV, Ayala V, Jove M, et al. Oxidative and endoplasmic reticulum stress interplay in sporadic amyotrophic lateral sclerosis. *Brain* 2007;130:3111-23.
300. Murata T, Ohtsuka C, Terayama Y. Increased mitochondrial oxidative damage in patients with sporadic amyotrophic lateral sclerosis. *J Neurol Sci* 2008;267:66-9.
301. Juhn SK, Ward WD. Alteration of oxidative enzymes (LDH and MDH) in perilymph after noise exposure. *Arch Otorhinolaryngol* 1979;222:103-8.
302. Van Campen LE, Murphy WJ, Franks JR, Mathias PI, Toraason MA. Oxidative DNA damage is associated with intense noise exposure in the rat. *Hear Res* 2002;164:29-38.
303. Bai U, Seidman MD, Hinojosa R, Quirk WS. Mitochondrial DNA deletions associated with aging and possibly presbycusis: a human archival temporal bone study. *Am J Otol* 1997;18:449-53.
304. Manwaring N, Jones MM, Wang JJ, et al. Mitochondrial DNA haplogroups and age-related hearing loss. *Arch Otolaryngol Head Neck Surg* 2007;133:929-33.
305. Seidman MD, Ahmad N, Joshi D, Seidman J, Thawani S, Quirk WS. Age-related hearing loss and its association with reactive oxygen species and mitochondrial DNA damage. *Acta Otolaryngol Suppl* 2004;16-24.
306. Seidman MD, Khan MJ, Bai U, Shirwany N, Quirk WS. Biologic activity of mitochondrial metabolites on aging and age-related hearing loss. *Am J Otol* 2000;21:161-7.
307. Yamasoba T, Someya S, Yamada C, Weindruch R, Prolla TA, Tanokura M. Role of mitochondrial dysfunction and mitochondrial DNA mutations in age-related hearing loss. *Hear Res* 2007;226:185-93.
308. Compston JE, Vedi S, Stephen AB, et al. Reduced bone formation in UK Gulf War veterans: a bone histomorphometric study. *J Clin Pathol* 2002;55:897-9.
309. Gray GC, Smith TC, Kang HK, Knoke JD. Are Gulf War veterans suffering war-related illnesses? Federal and civilian hospitalizations examined, June 1991 to December 1994. *Am J Epidemiol* 2000;151:63-71.
310. Ozgocmen S, Kaya H, Fadillioglu E, Aydogan R, Yilmaz Z. Role of antioxidant systems, lipid peroxidation, and nitric oxide in postmenopausal osteoporosis. *Mol Cell Biochem* 2006.

311. Varanasi SS, Francis RM, Berger CE, Papiha SS, Datta HK. Mitochondrial DNA deletion associated oxidative stress and severe male osteoporosis. *Osteoporos Int* 1999;10:143-9.
312. Ochoa JJ, Quiles JL, Planells E, et al. Lifelong supplementation with coenzyme Q10 protects from bone mineral density loss in rats during aging. *Fourth Conference of the International Coenzyme Q10 Association, Conference Proceedings 2005;Los Angeles:149-50.*
313. Compston JE, Vedi S, Stephen AB, et al. Reduced bone formation after exposure to organophosphates. *Lancet* 1999;354:1791-2.
314. Elverland HH, Torbergson T. Audiologic findings in a family with mitochondrial disorder. *Am J Otol* 1991;12:459-65.
315. Blondon H, Polivka M, Joly F, Flourie B, Mikol J, Messing B. Digestive smooth muscle mitochondrial myopathy in patients with mitochondrial-neuro-gastro-intestinal encephalomyopathy (MNGIE). *Gastroenterol Clin Biol* 2005;29:773-8.
316. Guillausseau PJ, Massin P, Dubois-LaFargue D, et al. Maternally inherited diabetes and deafness: a multicenter study. *Ann Intern Med* 2001;134:721-8.
317. Hutchison WM, Thyagarajan D, Poulton J, et al. Clinical and molecular features of encephalomyopathy due to the A3302G mutation in the mitochondrial tRNA(Leu(UUR)) gene. *Arch Neurol* 2005;62:1920-3.
318. Suzuki S, Hinokio Y, Ohtomo M, et al. The effects of coenzyme Q10 treatment on maternally inherited diabetes mellitus and deafness, and mitochondrial DNA 3243 (A to G) mutation. *Diabetologia* 1998;41:584-8.
319. Tanji K, Gamez J, Cervera C, et al. The A8344G mutation in mitochondrial DNA associated with stroke-like episodes and gastrointestinal dysfunction. *Acta Neuropathol (Berl)* 2003;105:69-75.
320. Wortmann S, Rodenburg RJ, Huizing M, et al. Association of 3-methylglutaconic aciduria with sensori-neural deafness, encephalopathy, and Leigh-like syndrome (MEGDEL association) in four patients with a disorder of the oxidative phosphorylation. *Mol Genet Metab* 2006.
321. Wray SH, Provenzale JM, Johns DR, Thulborn KR. MR of the brain in mitochondrial myopathy. *AJNR Am J Neuroradiol* 1995;16:1167-73.
322. Yuan H, Qian Y, Xu Y, et al. Cosegregation of the G7444A mutation in the mitochondrial COI/tRNA(Ser(UCN)) genes with the 12S rRNA A1555G mutation in a Chinese family with aminoglycoside-induced and nonsyndromic hearing loss. *Am J Med Genet A* 2005;138A:133-40.
323. Fechter LD, Gearhart C, Fulton S, et al. JP-8 Jet Fuel Can Promote Auditory Impairment Resulting From Subsequent Noise Exposure in Rats. *Toxicol Sci* 2007.
324. Nicotera TM, Ding D, McFadden SL, Salvemini D, Salvi R. Paraquat-induced hair cell damage and protection with the superoxide dismutase mimetic m40403. *Audiol Neurotol* 2004;9:353-62.
325. Dobrian AD, Davies MJ, Schriver SD, Lauterio TJ, Prewitt RL. Oxidative stress in a rat model of obesity-induced hypertension. *Hypertension* 2001;37:554-60.
326. Cottone S, Mule G, Nardi E, et al. Relation of C-reactive protein to oxidative stress and to endothelial activation in essential hypertension. *Am J Hypertens* 2006;19:313-8.
327. Vaziri ND. Pathogenesis of lead-induced hypertension: role of oxidative stress. *J Hypertens Suppl* 2002;20:S15-20.
328. Vaziri ND, Sica DA. Lead-induced hypertension: role of oxidative stress. *Curr Hypertens Rep* 2004;6:314-20.
329. Allen T, Rana SV. Oxidative stress by inorganic arsenic: modulation by thyroid hormones in rat. *Comp Biochem Physiol C Toxicol Pharmacol* 2003;135:157-62.
330. Kwok RK, Mendola P, Liu ZY, et al. Drinking water arsenic exposure and blood pressure in healthy women of reproductive age in Inner Mongolia, China. *Toxicol Appl Pharmacol* 2007;222:337-43.
331. Huang YK, Tseng CH, Huang YL, Yang MH, Chen CJ, Hsueh YM. Arsenic methylation capability and hypertension risk in subjects living in arseniasis-hyperendemic areas in southwestern Taiwan. *Toxicol Appl Pharmacol* 2007;218:135-42.

332. Borthwick GM, Johnson MA, Ince PG, Shaw PJ, Turnbull DM. Mitochondrial enzyme activity in amyotrophic lateral sclerosis: implications for the role of mitochondria in neuronal cell death. *Ann Neurol* 1999;46:787-90.
333. Aguirre N, Beal MF, Matson WR, Bogdanov MB. Increased oxidative damage to DNA in an animal model of amyotrophic lateral sclerosis. *Free Radic Res* 2005;39:383-8.
334. Albers DS, Beal MF. Mitochondrial dysfunction and oxidative stress in aging and neurodegenerative disease. *J Neural Transm Suppl* 2000;59:133-54.
335. Barber SC, Mead RJ, Shaw PJ. Oxidative stress in ALS: A mechanism of neurodegeneration and a therapeutic target. *Biochim Biophys Acta* 2006.
336. Mancuso M, Coppede F, Migliore L, Siciliano G, Murri L. Mitochondrial dysfunction, oxidative stress and neurodegeneration. *J Alzheimers Dis* 2006;10:59-73.
337. Liu R, Li B, Flanagan SW, Oberley LW, Gozal D, Qiu M. Increased mitochondrial antioxidative activity or decreased oxygen free radical propagation prevent mutant SOD1-mediated motor neuron cell death and increase amyotrophic lateral sclerosis-like transgenic mouse survival. *J Neurochem* 2002;80:488-500.
338. Mitsumoto H, Santella RM, Liu X, et al. Oxidative stress biomarkers in sporadic ALS. *Amyotroph Lateral Scler* 2008;9:177-83.
339. Murata T, Ohtsuka C, Terayama Y. Increased mitochondrial oxidative damage and oxidative DNA damage contributes to the neurodegenerative process in sporadic amyotrophic lateral sclerosis. *Free Radic Res* 2008;42:221-5.
340. Guegan C, Vila M, Rosoklija G, Hays AP, Przedborski S. Recruitment of the mitochondrial-dependent apoptotic pathway in amyotrophic lateral sclerosis. *J Neurosci* 2001;21:6569-76.
341. Kikuchi H, Furuta A, Nishioka K, Suzuki SO, Nakabeppu Y, Iwaki T. Impairment of mitochondrial DNA repair enzymes against accumulation of 8-oxo-guanine in the spinal motor neurons of amyotrophic lateral sclerosis. *Acta Neuropathol (Berl)* 2002;103:408-14.
342. Kong J, Xu Z. Massive mitochondrial degeneration in motor neurons triggers the onset of amyotrophic lateral sclerosis in mice expressing a mutant SOD1. *J Neurosci* 1998;18:3241-50.
343. Wiedemann FR, Manfredi G, Mawrin C, Beal MF, Schon EA. Mitochondrial DNA and respiratory chain function in spinal cords of ALS patients. *J Neurochem* 2002;80:616-25.
344. Cronin S, Greenway MJ, Prehn JH, Hardiman O. Paraoxonase promoter and intronic variants modify risk of sporadic amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 2007;78:984-6.
345. Morahan JM, Yu B, Trent RJ, Pamphlett R. A gene-environment study of the paraoxonase 1 gene and pesticides in amyotrophic lateral sclerosis. *Neurotoxicology* 2007;28:532-40.
346. Slowik A, Tomik B, Partyka D, et al. Paraoxonase-1 Q192R polymorphism and risk of sporadic amyotrophic lateral sclerosis. *Clin Genet* 2006;69:358-9.
347. Saeed M, Siddique N, Hung WY, et al. Paraoxonase cluster polymorphisms are associated with sporadic ALS. *Neurology* 2006.
348. La Du BN, Billecke S, Hsu C, Haley RW, Broomfield CA. Serum paraoxonase (PON1) isozymes: the quantitative analysis of isozymes affecting individual sensitivity to environmental chemicals. *Drug Metab Dispos* 2001;29:566-9.
349. Mackness B, Durrington P, Povey A, et al. Paraoxonase and susceptibility to organophosphorus poisoning in farmers dipping sheep. *Pharmacogenetics* 2003;13:81-8.
350. Povey AC, Mackness MI, Durrington PN, et al. Paraoxonase polymorphisms and self-reported chronic ill-health in farmers dipping sheep. *Occup Med (Lond)* 2005;55:282-6.
351. Golomb BA, Kwon EK, Koperski S, Evans MA. Amyotrophic lateral sclerosis-like conditions in possible association with cholesterol-lowering drugs: an analysis of patient reports to the University of California, San Diego (UCSD) Statin Effects Study. *Drug Saf* 2009;32:649-61.
352. Dupuis L, di Scala F, Rene F, et al. Up-regulation of mitochondrial uncoupling protein 3 reveals an early muscular metabolic defect in amyotrophic lateral sclerosis. *Faseb J* 2003;17:2091-3.
353. Dupuis L, Gonzalez de Aguilar JL, Echaniz-Laguna A, Loeffler JP. Mitochondrial dysfunction in amyotrophic lateral sclerosis also affects skeletal muscle. *Muscle Nerve* 2006;34:253-4.

354. Krasnianski A, Deschauer M, Neudecker S, et al. Mitochondrial changes in skeletal muscle in amyotrophic lateral sclerosis and other neurogenic atrophies. *Brain* 2005;128:1870-6.
355. Vielhaber S, Kornblum C, Heinze HJ, Elger CE, Kunz WS. Mitochondrial changes in skeletal muscle in amyotrophic lateral sclerosis and other neurogenic atrophies--a comment. *Brain* 2005;128:E38.
356. Vielhaber S, Kudin A, Winkler K, et al. Is there mitochondrial dysfunction in amyotrophic lateral sclerosis skeletal muscle? *Ann Neurol* 2003;53:686-7; author reply 7-8.
357. Echaniz-Laguna A, Zoll J, Ponsot E, et al. Muscular mitochondrial function in amyotrophic lateral sclerosis is progressively altered as the disease develops: a temporal study in man. *Exp Neurol* 2006;198:25-30.
358. Ringholz GM, Appel SH, Bradshaw M, Cooke NA, Mosnik DM, Schulz PE. Prevalence and patterns of cognitive impairment in sporadic ALS. *Neurology* 2005;65:586-90.
359. Wheaton MW, Salamone AR, Mosnik DM, et al. Cognitive impairment in familial ALS. *Neurology* 2007;69:1411-7.
360. Miele L, Gabrieli ML, Forgione A, et al. [Oxidative stress in metabolic syndrome and nonalcoholic steatohepatitis. Is it possible a role for vitamins in clinical practice?]. *Recenti Prog Med* 2006;97:1-5.
361. Grattagliano I, Palmieri VO, Portincasa P, Moschetta A, Palasciano G. Oxidative stress-induced risk factors associated with the metabolic syndrome: a unifying hypothesis. *J Nutr Biochem* 2008;19:491-504.
362. Williams TA, Mars AE, Buyske SG, et al. Risk of autistic disorder in affected offspring of mothers with a glutathione S-transferase P1 haplotype. *Arch Pediatr Adolesc Med* 2007;161:356-61.
363. Abd El-Gawad HM, El-Sawalhi MM. Nitric oxide and oxidative stress in brain and heart of normal rats treated with doxorubicin: role of aminoguanidine. *J Biochem Mol Toxicol* 2004;18:69-77.
364. Abu-Qare AW, Abou-Donia MB. Combined exposure to DEET (N,N-diethyl-m-toluamide) and permethrin: pharmacokinetics and toxicological effects. *J Toxicol Environ Health B Crit Rev* 2003;6:41-53.
365. Abushamaa AM, Sporn TA, Folz RJ. Oxidative stress and inflammation contribute to lung toxicity after a common breast cancer chemotherapy regimen. *Am J Physiol Lung Cell Mol Physiol* 2002;283:L336-45.
366. Bandyopadhyay D, Ghosh G, Bandyopadhyay A, Reiter RJ. Melatonin protects against piroxicam-induced gastric ulceration. *J Pineal Res* 2004;36:195-203.
367. Basivireddy J, Jacob M, Balasubramanian KA. Indomethacin induces free radical-mediated changes in renal brush border membranes. *Arch Toxicol* 2005;79:441-50.
368. Basivireddy J, Jacob M, Pulimood AB, Balasubramanian KA. Indomethacin-induced renal damage: role of oxygen free radicals. *Biochem Pharmacol* 2004;67:587-99.
369. Boelsterli UA. Mechanisms of NSAID-induced hepatotoxicity: focus on nimesulide. *Drug Saf* 2002;25:633-48.
370. Conklin KA. Coenzyme q10 for prevention of anthracycline-induced cardiotoxicity. *Integr Cancer Ther* 2005;4:110-30.
371. Denicola A, Radi R. Peroxynitrite and drug-dependent toxicity. *Toxicology* 2005;208:273-88.
372. Goli AK, Goli SA, Byrd RP, Jr., Roy TM. Simvastatin-induced lactic acidosis: a rare adverse reaction? *Clin Pharmacol Ther* 2002;72:461-4.
373. Hawkey CJ. Non-steroidal anti-inflammatory drug gastropathy: causes and treatment. *Scand J Gastroenterol Suppl* 1996;220:124-7.
374. Maldonado PD, Barrera D, Rivero I, et al. Antioxidant S-allylcysteine prevents gentamicin-induced oxidative stress and renal damage. *Free Radic Biol Med* 2003;35:317-24.
375. Naidu PS, Singh A, Kulkarni SK. Quercetin, a bioflavonoid, attenuates haloperidol-induced orofacial dyskinesia. *Neuropharmacology* 2003;44:1100-6.
376. Perez de Hornedo J, de Arriba G, Calvino M, Benito S, Parra T. [Cyclosporin A causes oxidative stress and mitochondrial dysfunction in renal tubular cells]. *Nefrologia* 2007;27:565-73.
377. Weyers AI, Ugnia LI, Garcia Ovando H, Gorla NB. Ciprofloxacin increases hepatic and renal lipid hydroperoxides levels in mice. *Biocell* 2002;26:225-8.

378. Carbonera D, Angrilli A, Azzone GF. Mechanism of nitrofurantoin toxicity and oxidative stress in mitochondria. *Biochim Biophys Acta* 1988;936:139-47.
379. Poitras P, Gougeon A, Binn M, Bouin M. Extra digestive manifestations of irritable bowel syndrome: intolerance to drugs? *Dig Dis Sci* 2008;53:2168-76.
380. Ashfaq S, Abramson JL, Jones DP, et al. Endothelial function and aminothiols biomarkers of oxidative stress in healthy adults. *Hypertension* 2008;52:80-5.
381. Mitsuyoshi H, Itoh Y, Okanoue T. [Role of oxidative stress in non-alcoholic steatohepatitis]. *Nippon Rinsho* 2006;64:1077-82.
382. Natarajan SK, Eapen CE, Pullimood AB, Balasubramanian KA. Oxidative stress in experimental liver microvesicular steatosis: role of mitochondria and peroxisomes. *J Gastroenterol Hepatol* 2006;21:1240-9.
383. Grattagliano I, Palmieri VO, Portincasa P, Moschetta A, Palasciano G. Oxidative stress-induced risk factors associated with the metabolic syndrome: a unifying hypothesis. *J Nutr Biochem* 2007.
384. Tripathy D, Aljada A, Dandona P. Free fatty acids (FFA) and endothelial dysfunction; role of increased oxidative stress and inflammation. --to: Steinberg et al. (2002) Vascular function, insulin resistance and fatty acids. *Diabetologia* 2003;46:300-1.
385. Pereira EC, Ferderbar S, Bertolami MC, et al. Biomarkers of oxidative stress and endothelial dysfunction in glucose intolerance and diabetes mellitus. *Clin Biochem* 2008;41:1454-60.
386. Lamson DW, Plaza SM. Mitochondrial factors in the pathogenesis of diabetes: a hypothesis for treatment. *Altern Med Rev* 2002;7:94-111.
387. Belch JJ, Mackay IR, Hill A, Jennings P, McCollum P. Oxidative stress is present in atherosclerotic peripheral arterial disease and further increased by diabetes mellitus. *Int Angiol* 1995;14:385-8.
388. Loffredo L, Marcoccia A, Pignatelli P, et al. Oxidative-stress-mediated arterial dysfunction in patients with peripheral arterial disease. *Eur Heart J* 2007;28:608-12.
389. Mueller T, Dieplinger B, Gegenhuber A, et al. Serum total 8-iso-prostaglandin F2alpha: a new and independent predictor of peripheral arterial disease. *J Vasc Surg* 2004;40:768-73.
390. Bhat HK, Hiatt WR, Hoppel CL, Brass EP. Skeletal muscle mitochondrial DNA injury in patients with unilateral peripheral arterial disease. *Circulation* 1999;99:807-12.
391. Hiatt WR. Carnitine and peripheral arterial disease. *Ann N Y Acad Sci* 2004;1033:92-8.
392. Makris KI, Nella AA, Zhu Z, et al. Mitochondriopathy of peripheral arterial disease. *Vascular* 2007;15:336-43.
393. Shoffner JM, Watts RL, Juncos JL, Torroni A, Wallace DC. Mitochondrial oxidative phosphorylation defects in Parkinson's disease. *Ann Neurol* 1991;30:332-9.
394. Haas RH, Nasirian F, Nakano K, et al. Low platelet mitochondrial complex I and complex II/III activity in early untreated Parkinson's disease. *Ann Neurol* 1995;37:714-22.
395. Benmoyal-Segal L, Vander T, Shifman S, et al. Acetylcholinesterase/paraoxonase interactions increase the risk of insecticide-induced Parkinson's disease. *Faseb J* 2005;19:452-4.
396. Zintzaras E, Hadjigeorgiou GM. Association of paraoxonase 1 gene polymorphisms with risk of Parkinson's disease: a meta-analysis. *J Hum Genet* 2004;49:474-81.
397. Carmine A, Buervenich S, Sydow O, Anvret M, Olson L. Further evidence for an association of the paraoxonase 1 (PON1) Met-54 allele with Parkinson's disease. *Mov Disord* 2002;17:764-6.
398. Akhmedova SN, Yakimovsky AK, Schwartz EI. Paraoxonase 1 Met--Leu 54 polymorphism is associated with Parkinson's disease. *J Neurol Sci* 2001;184:179-82.
399. Kondo I, Yamamoto M. Genetic polymorphism of paraoxonase 1 (PON1) and susceptibility to Parkinson's disease. *Brain Res* 1998;806:271-3.
400. Gardner A, Boles RG. Symptoms of somatization as a rapid screening tool for mitochondrial dysfunction in depression. *Biopsychosoc Med* 2008;2:7.
401. Bullman TA, Mahan CM, Kang HK, Page WF. Mortality in US Army Gulf War veterans exposed to 1991 Khamisiyah chemical munitions destruction. *Am J Public Health* 2005;95:1382-8.

402. Barth SK, Kang HK, Bullman TA, Wallin MT. Neurological mortality among U.S. veterans of the Persian Gulf War: 13-year follow-up. Am J Ind Med 2009;52:663-70.